

Rabbit anti-Rat SECTM1 Polyclonal Antibody | anti-SECTM1 antibody

SECTM1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291274	Rabbit	Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

SECTM1; Polyclonal Antibody; SECTM1 Polyclonal Antibody; Secreted and transmembrane protein 1; Protein K-12; K12; anti-SECTM1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-SECTM1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (SECTM1) Polyclonal Antibody, Unconjugated (bs-10153R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (pv-0023) instructions and DAB staining.




IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (SECTM1) Polyclonal Antibody, Unconjugated (bs-10153R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (rat live); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (SECTM1) Polyclonal Antibody, Unconjugated (bs-10153R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37 degree C for 20 min; Incubation: Anti-SECTM1 Polyclonal Antibody, Unconjugated (bs-10153R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506869
NCBI GeneID	6398

NCBI Accession #	NP_002995.1
NCBI GenBank Nucleotide #	NM_003004.2
UniProt Accession #	Q8WVN6 ; O00466 ; B2R7H0
Molecular Weight	27,039 Da
NCBI Official Full Name	secreted and transmembrane protein 1
NCBI Official Synonym Full Names	secreted and transmembrane 1
NCBI Official Symbol	SECTM1
NCBI Official Synonym Symbols	K12
NCBI Protein Information	secreted and transmembrane protein 1
UniProt Protein Name	Secreted and transmembrane protein 1
UniProt Gene Name	SECTM1
UniProt Synonym Gene Names	K12

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/sectm1-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291274>