

Rabbit anti-Rat, Cow NR5A2 Polyclonal Antibody | anti-NR5A2 antibody

NR5A2 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291282	Rabbit	Rat, Cow	Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

NR5A2; Polyclonal Antibody; NR5A2 Polyclonal Antibody; Nuclear receptor subfamily 5 group A member 2; Alpha-1-fetoprotein transcription factor; B1-binding factor; hB1F; CYP7A promoter-binding factor; Hepatocytic transcription factor; Liver receptor homolog 1; LRH-1; B1F; CPF; FTF; anti-NR5A2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Rat, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-NR5A2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




WB (Western Blot)

Dilution: Sample:Liver(Rat) Lysate at 40 ugPrimary: Anti-NR5A2 (bs-10444R) at 1/1000
 dilutionSecondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilutionPredicted band size:
 61 kDObserved band size: 61 kD



IHC (Immunohistochemistry)

Tissue/cell: mouse embryo tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min;Incubation: Anti-NR5A2 Polyclonal Antibody, Unconjugated(bs-10444R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (NR5A2) Polyclonal Antibody, Unconjugated (bs-10444R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	451172094
NCBI GeneID	2494

NCBI Accession #	NP_001263393.1
NCBI GenBank Nucleotide #	NM_001276464.1
UniProt Accession #	O00482; O95642; Q147U3; B4E2P3
Molecular Weight	53,657 Da
NCBI Official Full Name	nuclear receptor subfamily 5 group A member 2 isoform 3
NCBI Official Synonym Full Names	nuclear receptor subfamily 5 group A member 2
NCBI Official Symbol	NR5A2
NCBI Official Synonym Symbols	B1F; CPF; FTF; B1F2; LRH1; LRH-1; FTZ-F1; hB1F-2; FTZ-F1beta
NCBI Protein Information	nuclear receptor subfamily 5 group A member 2
UniProt Protein Name	Nuclear receptor subfamily 5 group A member 2
UniProt Gene Name	NR5A2
UniProt Synonym Gene Names	B1F; CPF; FTF; hB1F; LRH-1
UniProt Entry Name	NR5A2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/nr5a2-antibody-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291282>