

Rabbit TIMP-1 Polyclonal Antibody | anti-TIMP-1 antibody

TIMP-1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA29132	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunofluorescence

PRODUCT SYNONYMS

TIMP-1; Polyclonal Antibody; TIMP-1 Polyclonal Antibody; TIMP1; CLGI; TIMP; Metalloproteinase inhibitor 1; Erythroid-potentiating activity; EPA; Fibroblast collagenase inhibitor; Collagenase inhibitor; Tissue inhibitor of metalloproteinases 1; anti-TIMP-1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Applicable Applications for anti-TIMP-1 antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence)
Application Notes	IF: 1:50-200 WB: 1/500-1/2000 ELISA: 1/10000
Cellular Location	Secreted
Tissue Location	Detected in rheumatoid synovial fluid (at protein level).
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.



IHC (Immunohistochemistry)

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.



IF (Immunofluorescence)

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.

**Application Data**

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.

**Application Data**

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-TIMP-1 antibody

Metalloproteinase inhibitor that functions by forming one to one complexes with target metalloproteinases, such as collagenases, and irreversibly inactivates them by binding to their catalytic zinc cofactor. Acts on MMP1, MMP2, MMP3, MMP7, MMP8, MMP9, MMP10, MMP11, MMP12, MMP13 and MMP16. Does not act on MMP14. Also functions as a growth factor that regulates cell differentiation, migration and cell death and activates cellular signaling cascades via CD63 and ITGB1. Plays a role in integrin signaling. Mediates erythropoiesis in vitro; but, unlike IL3, it is species-specific, stimulating the growth and differentiation of only human and murine erythroid progenitors.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507509
NCBI GeneID	7076
NCBI Accession #	NP_003245.1
NCBI GenBank Nucleotide #	NM_003254.2
UniProt Accession #	P01033; Q14252; Q9UCU1
Molecular Weight	23,171 Da
NCBI Official Full Name	metalloproteinase inhibitor 1
NCBI Official Synonym Full Names	TIMP metalloproteinase inhibitor 1
NCBI Official Symbol	TIMP1
NCBI Official Synonym Symbols	EPA; EPO; HCI; CLGI; TIMP
NCBI Protein Information	metalloproteinase inhibitor 1; TIMP-1; collagenase inhibitor; erythroid potentiating activity; erythroid-potentiating activity; fibroblast collagenase inhibitor; tissue inhibitor of metalloproteinases 1
UniProt Protein Name	Metalloproteinase inhibitor 1
UniProt Gene Name	TIMP1
UniProt Synonym Gene Names	CLGI; TIMP; EPA; Collagenase inhibitor; TIMP-1
UniProt Entry Name	TIMP1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/timp-1-antibody-western-blot-immunohistochemistry-immunofluorescence-elisa/29132>