

Rabbit VEGF-D Polyclonal Antibody | anti-VEGF-D antibody

VEGF-D Polyclonal Antibody

CATALOG # AAA29136	HOST Rabbit	REACTIVITY Human, Mouse, Rat, Monkey	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	--	---


PRODUCT SYNONYMS


VEGF-D; Polyclonal Antibody; VEGF-D Polyclonal Antibody; FIGF; VEGFD; Vascular endothelial growth factor D; c-Fos-induced growth factor; anti-VEGF-D antibody


PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Applicable Applications for anti-VEGF-D antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.

IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.

IHC (Immunohistochemistry)


Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.

IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.
Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-VEGF-D antibody

Growth factor active in angiogenesis, lymphangiogenesis and endothelial cell growth, stimulating their proliferation and migration and also has effects on the permeability of blood vessels. May function in the formation of the venous and lymphatic vascular systems during embryogenesis, and also in the maintenance of differentiated lymphatic endothelium in adults. Binds and activates VEGFR-2 (KDR/FLK1) and VEGFR-3 (FLT4) receptors.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758378
NCBI GeneID	2277
NCBI Accession #	NP_004460.1
NCBI GenBank Nucleotide #	NM_004469.4
UniProt Accession #	O43915; B2R7Z3
Molecular Weight	40,444 Da
NCBI Official Full Name	vascular endothelial growth factor D preproprotein
NCBI Official Synonym Full Names	c-fos induced growth factor (vascular endothelial growth factor D)
NCBI Official Symbol	FIGF
NCBI Official Synonym Symbols	VEGFD; VEGF-D
NCBI Protein Information	vascular endothelial growth factor D
UniProt Protein Name	Vascular endothelial growth factor D
UniProt Gene Name	FIGF
UniProt Synonym Gene Names	VEGFD; VEGF-D; FIGF
UniProt Entry Name	VEGFD_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/vegfd-antibody-western-blot-immunohistochemistry-elisa/29136>