

Rabbit CRBN Polyclonal Antibody | anti-CRBN antibody

CRBN Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291360	Rabbit	Rat, Pig, Dog, Cow	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry


PRODUCT SYNONYMS

CRBN; Polyclonal Antibody; CRBN Polyclonal Antibody; Protein cereblon; anti-CRBN antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Pig, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-CRBN antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (CRBN) Polyclonal Antibody, Unconjugated (bs-11716R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37 degree C for 20 min; Incubation: Anti-CRBN Polyclonal Antibody, Unconjugated (bs-11716R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



IHC (Immunohistochemistry)

Tissue/cell: rat testis tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37 degree C for 20 min; Incubation: Anti-CRBN Polyclonal Antibody, Unconjugated (bs-11716R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining

NCBI AND UNIPROT INFORMATION

NCBI GI #	39545580
NCBI GeneID	51185

NCBI Accession #	NP_057386
NCBI GenBank Nucleotide #	NM_016302.3
UniProt Accession #	Q96SW2; Q6AI62; Q6NVZ0; Q9UHW4; B2R6H4; C9IZA9; C9JAH6
Molecular Weight	Predicted: 49 kDa Observed: 54 kDa
NCBI Official Full Name	protein cereblon isoform 1
NCBI Official Synonym Full Names	cereblon
NCBI Official Symbol	CRBN
NCBI Official Synonym Symbols	MRT2; MRT2A
NCBI Protein Information	protein cereblon; protein x 0001
UniProt Protein Name	Protein cereblon
UniProt Gene Name	CRBN
UniProt Entry Name	CRBN_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/crbn-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-elisa/291360>