

Rabbit anti-Rat, Dog CSAD Polyclonal Antibody | anti-CSAD antibody

CSAD Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291368	Rabbit	Rat, Dog	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CSAD; Polyclonal Antibody; CSAD Polyclonal Antibody; Cysteine sulfinic acid decarboxylase; 4.1.1.29; Cysteine-sulfinate decarboxylase; Sulfinoalanine decarboxylase; CSD; anti-CSAD antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Rat, Dog
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-CSAD antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (CSAD) Polyclonal Antibody, Unconjugated (bs-11822R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (CSAD) Polyclonal Antibody, Unconjugated (bs-11822R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	347659021
NCBI GeneID	51380
NCBI Accession #	NP_001231634.1
NCBI GenBank Nucleotide #	NM_001244705.1
UniProt Accession #	Q9Y600; Q4QQH9; Q9UNJ5; Q9Y601; A8K0U4
Molecular Weight	55,023 Da
NCBI Official Full Name	cysteine sulfinic acid decarboxylase isoform 2
NCBI Official Synonym Full Names	cysteine sulfinic acid decarboxylase

NCBI Official Symbol	CSAD
NCBI Official Synonym Symbols	CSD; PCAP
NCBI Protein Information	cysteine sulfinic acid decarboxylase; sulfinoalanine decarboxylase; cysteine-sulfinate decarboxylase; P-selectin cytoplasmic tail-associated protein; cysteine sulfinic acid decarboxylase-related protein
UniProt Protein Name	Cysteine sulfinic acid decarboxylase
UniProt Gene Name	CSAD
UniProt Synonym Gene Names	CSD
UniProt Entry Name	CSAD_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/csad-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291368>