

Rabbit XIAP Polyclonal Antibody | anti-XIAP antibody

XIAP Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA29138	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry


PRODUCT SYNONYMS

XIAP; Polyclonal Antibody; XIAP Polyclonal Antibody; API3; BIRC4; IAP3; E3 ubiquitin-protein ligase XIAP; Baculoviral IAP repeat-containing protein 4; IAP-like protein; ILP; hILP; Inhibitor of apoptosis protein 3; IAP-3; hIAP-3; hIAP3; X-linked inhibitor of apoptosis protein; X-linked I; anti-XIAP antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-XIAP antibody	WB (Western Blot), IHC (Immunohistochemistry)
Cellular Location	Cytoplasm. Nucleus. Note=TLE3 promotes its nuclear localization
Tissue Location	Ubiquitous, except peripheral blood leukocytes.
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.
Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.
Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.
Not yet tested in other applications.




IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.
Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.
Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000.
Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-XIAP antibody

Multi-functional protein which regulates not only caspases and apoptosis, but also modulates inflammatory signaling and immunity, copper homeostasis, mitogenic kinase signaling, cell proliferation, as well as cell invasion and metastasis. Acts as a direct caspase inhibitor. Directly bind to the active site pocket of CASP3 and CASP7 and obstructs substrate entry. Inactivates CASP9 by keeping it in a monomeric, inactive state. Acts as an E3 ubiquitin-protein ligase regulating NF-kappa-B signaling and the target proteins for its E3 ubiquitin-protein ligase activity include: RIPK1, CASP3, CASP7, CASP8, CASP9, MAP3K2/MEKK2, DIABLO/SMAC, AIFM1, CCS and BIRC5/survivin. Ubiquitination of CCS leads to enhancement of its chaperone activity toward its physiologic target, SOD1, rather than proteasomal degradation. Ubiquitination of MAP3K2/MEKK2 and AIFM1 does not lead to proteasomal degradation. Plays a role in copper homeostasis by ubiquitinating COMMD1 and promoting its proteasomal degradation. Can also function as E3 ubiquitin-protein ligase of the NEDD8 conjugation pathway, targeting effector caspases for neddylation and inactivation. Regulates the BMP signaling pathway and the SMAD and MAP3K7/TAK1 dependent pathways leading to NF-kappa-B and JNK activation. Acts as an important regulator of innate immune signaling via regulation of Nodlike receptors (NLRs). Protects cells from spontaneous formation of the ripoptosome, a large multi-protein complex that has the capability to kill cancer cells in a caspase-dependent and caspase-independent manner. Suppresses ripoptosome formation by ubiquitinating RIPK1 and CASP8. Acts as a positive regulator of Wnt signaling and ubiquitinates TLE1, TLE2, TLE3, TLE4 and AES. Ubiquitination of TLE3 results in inhibition of its interaction with TCF7L2/TCF4 thereby allowing efficient recruitment and binding of the transcriptional coactivator beta- catenin to TCF7L2/TCF4 that is required to initiate a Wnt-specific transcriptional program.

NCBI AND UNIPROT INFORMATION

NCBI GI #	32528299
NCBI GeneID	331
NCBI Accession #	NP_001158.2

NCBI GenBank Nucleotide #	NM_001167.3
UniProt Accession #	P98170 ; Q9NQ14 ; D3DTF2
Molecular Weight	56,685 Da
NCBI Official Full Name	E3 ubiquitin-protein ligase XIAP
NCBI Official Synonym Full Names	X-linked inhibitor of apoptosis
NCBI Official Symbol	XIAP
NCBI Official Synonym Symbols	API3; ILP1; MIHA; XLP2; BIRC4; IAP-3; hIAP3; hIAP-3
NCBI Protein Information	E3 ubiquitin-protein ligase XIAP; X-linked IAP; IAP-like protein; inhibitor of apoptosis protein 3; baculoviral IAP repeat-containing protein 4
UniProt Protein Name	E3 ubiquitin-protein ligase XIAP
UniProt Gene Name	XIAP
UniProt Synonym Gene Names	API3; BIRC4; IAP3; ILP; hILP; IAP-3; hIAP-3; hIAP3
UniProt Entry Name	XIAP_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/xiap-antibody-western-blot-immunohistochemistry-elisa/29138>