

Rabbit anti-Rat TUG Polyclonal Antibody | anti-TUG antibody

TUG Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291402	Rabbit	Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

TUG; Polyclonal Antibody; TUG Polyclonal Antibody; Tether containing UBX domain for GLUT4; Alveolar soft part sarcoma chromosomal region candidate gene 1 protein; Alveolar soft part sarcoma locus; Renal papillary cell carcinoma protein 17; UBX domain-containing protein 9; ASPSCR1; ASPL; RCC17; UBXD9; UBXN9; anti-TUG antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-TUG antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (TUG) Polyclonal Antibody, Unconjugated (bs-12355R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (rat colon tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (TUG) Polyclonal Antibody, Unconjugated (bs-12355R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



WB (Western Blot)

Sample:Heart (Mouse) Lysate at 40 ugPrimary: Anti-TUG (bs-12355R) at 1/1000 dilutionSecondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilutionPredicted band size: 60 kDObserved band size: 58 kD

NCBI AND UNIPROT INFORMATION

NCBI GI #	353677961
NCBI GeneID	79058
NCBI Accession #	NP_001238817.1

NCBI GenBank Nucleotide #	NM_001251888.1
UniProt Accession #	Q9BZE9; Q7Z6N7; Q8WV59; Q96LS5; Q96M40; A8K3K9
Molecular Weight	37,910 Da
NCBI Official Full Name	tether containing UBX domain for GLUT4 isoform 2
NCBI Official Synonym Full Names	alveolar soft part sarcoma chromosome region, candidate 1
NCBI Official Symbol	ASPC1
NCBI Official Synonym Symbols	TUG; ASPL; ASPS; RCC17; UBXD9; UBXN9; ASPCR1
NCBI Protein Information	tether containing UBX domain for GLUT4; UBX domain protein 9; UBX domain-containing protein 9; alveolar soft part sarcoma chromosomal region candidate gene 1 protein; alveolar soft part sarcoma locus; renal cell carcinoma gene on chromosome 17; renal cell
UniProt Protein Name	Tether containing UBX domain for GLUT4
UniProt Gene Name	ASPC1
UniProt Synonym Gene Names	ASPL; RCC17; TUG; UBXD9; UBXN9
UniProt Entry Name	ASPC1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tug-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291402>