# Rabbit anti-Rat SNRNP200 Polyclonal Antibody | anti-SNRNP200 antibody

SNRNP200 Polyclonal Antibody

| CATALOG # | HOST | REACTIVITY | APPLICATIONS |
|---|---|---|---|
| **AAA291421** | **Rabbit** | **Rat** | **Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry** |


## PRODUCT SYNONYMS

SNRNP200; Polyclonal Antibody; SNRNP200 Polyclonal Antibody; U5 small nuclear ribonucleoprotein 200 kDa helicase; 3.6.4.13; Activating signal cointegrator 1 complex subunit 3-like 1; BRR2 homolog; U5 snRNP-specific 200 kDa protein; U5-200KD; ASCC3L1; HELIC2; KIAA0788; anti-SNRNP200 antibody


## PRODUCT SPECIFICATIONS

| | |
|---|---|
| **Host** | Rabbit |
| **Reactivity** | Rat |
| **Clonality** | Polyclonal |
| **Purity/Purification** | Purified Rabbit Polyclonal Antibody (Pab) |
| **Applicable Applications for anti-SNRNP200 antibody** | ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry) |
| **Preparation and Storage** | Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles. |

## IMAGES



**IHC (Immunohistochemisry)**
Dilution: Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (SNRNP200) Polyclonal Antibody, Unconjugated (bs-12657R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



**IHC (Immunohiostchemistry)**
Paraformaldehyde-fixed, paraffin embedded (Rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (SNRNP200) Polyclonal Antibody, Unconjugated (bs-12657R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



**FCM/FACS (Flow Cytometry)**
Blank control:U-937. Primary Antibody (green line): Rabbit Anti-SNRNP200 antibody (bs-12657R) Dilution: 2 ug /10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG .Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: 2 ug /test.Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at-20 degree C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

## NCBI AND UNIPROT INFORMATION

| | |
|---|---|
| **NCBI GI #** | 40217847 |

| | |
|---|---|
| **NCBI GeneID** | 23020 |
| **NCBI Accession #** | NP_054733.2 |
| **NCBI GenBank Nucleotide #** | NM_014014.4 |
| **UniProt Accession #** | O75643; O94884; Q6NZY0; Q6PX59; Q8NBE6; Q96IF2; Q9H7S0 |
| **Molecular Weight** | 71,472 Da |
| **NCBI Official Full Name** | U5 small nuclear ribonucleoprotein 200 kDa helicase |
| **NCBI Official Synonym Full Names** | small nuclear ribonucleoprotein 200kDa (U5) |
| **NCBI Official Symbol** | SNRNP200 |
| **NCBI Official Synonym Symbols** | BRR2; RP33; HELIC2; ASCC3L1; U5-200KD |
| **NCBI Protein Information** | U5 small nuclear ribonucleoprotein 200 kDa helicase |
| **UniProt Protein Name** | U5 small nuclear ribonucleoprotein 200 kDa helicase |
| **UniProt Gene Name** | **SNRNP200** |
| **UniProt Synonym Gene Names** | ASCC3L1; HELIC2; KIAA0788; U5-200KD |
| **UniProt Entry Name** | U520_HUMAN |

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

**AAA Biotech** | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/polyclonal/snrnp200-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-elisa/291421