

Rabbit STK32C Polyclonal Antibody | anti-STK32C antibody

STK32C Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291437	Rabbit	Rat, Pig, Dog	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

STK32C; Polyclonal Antibody; STK32C Polyclonal Antibody; Serine/threonine-protein kinase 32C; 2.7.11.1; PKE; Yet another novel kinase 3; YANK3; anti-STK32C antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Pig, Dog
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-STK32C antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Notes	ECO:0000312 EMBL: AAH45760.1
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (STK32C) Polyclonal Antibody, Unconjugated (bs-12833R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (STK32C) Polyclonal Antibody, Unconjugated (bs-12833R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



WB (Western Blot)

Sample: Molt-4(Human) Cell Lysate at 40 ugA549(Human) Cell Lysate at 40 ugPrimary: Anti-STK32C (bs-12833R) at 1/300 dilutionSecondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilutionPredicted band size: 56 kDobserved band size: 56 kD

NCBI AND UNIPROT INFORMATION

NCBI GI #	32455269
-----------	----------

NCBI GeneID	282974
NCBI Accession #	NP_775846.2
NCBI GenBank Nucleotide #	NM_173575.2
UniProt Accession #	Q86UX6 ; Q5T0Q5 ; Q86UE1
Molecular Weight	54,994 Da
NCBI Official Full Name	serine/threonine-protein kinase 32C
NCBI Official Synonym Full Names	serine/threonine kinase 32C
NCBI Official Symbol	STK32C
NCBI Official Synonym Symbols	PKE; YANK3; RP11-140A10.1
NCBI Protein Information	serine/threonine-protein kinase 32C; PKE protein kinase; protein kinase (PKE); yet another novel kinase 3
UniProt Protein Name	Serine/threonine-protein kinase 32C
UniProt Gene Name	STK32C
UniProt Synonym Gene Names	YANK3
UniProt Entry Name	ST32C_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/stk32c-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291437>