

Rabbit EXOSC10 Polyclonal Antibody | anti-EXOSC10 antibody

EXOSC10 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291450	Rabbit	Rat, Pig, Dog, Cow	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

EXOSC10; Polyclonal Antibody; EXOSC10 Polyclonal Antibody; Exosome component 10; 3.1.13.-; Autoantigen PM/Scl 2; P100 polymyositis-scleroderma overlap syndrome-associated autoantigen; Polymyositis/scleroderma autoantigen 100 kDa; PM/Scl-100; Polymyositis/scleroderma autoantigen 2; anti-EXOSC10 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Pig, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-EXOSC10 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (mouse skin); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (EXOSC10) Polyclonal Antibody, Unconjugated (bs-13120R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




IHC (Immunohistochemistry)

Dilution: Tissue/cell: Mouse Lymph nodes; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37 degree C for 20 min; Incubation: Anti-EXOSC10, Unconjugated (bs-13120R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



IHC (Immunohistochemistry)

Tissue/cell: human skin tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37 degree C for 20 min; Incubation: Anti-EXOSC10, Unconjugated (bs-13120R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



WB (Western Blot)

Sample: U937 (Human) Cell Lysate at 30 ug; Primary: Anti- EXOSC10 (bs-13120R) at 1/1000 dilution; Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution; Predicted band size: 101 kDa; Observed band size: 103 kDa

NCBI AND UNIPROT INFORMATION

NCBI GI #	8928564
-----------	---------

NCBI GeneID	5394
UniProt Accession #	Q01780; Q15158; B1AKQ0; B1AKQ1
Molecular Weight	885
NCBI Official Full Name	Exosome component 10
NCBI Official Synonym Full Names	exosome component 10
NCBI Official Symbol	EXOSC10
NCBI Official Synonym Symbols	p2; p3; p4; RRP6; PMSCL; Rrp6p; PM-Scl; PMSCL2; PM/Scl-100
NCBI Protein Information	exosome component 10; autoantigen PM-SCL; polymyositis/scleroderma autoantigen 2; polymyositis/scleroderma autoantigen 100 kDa; P100 polymyositis-scleroderma overlap syndrome-associated autoantigen
UniProt Protein Name	Exosome component 10
UniProt Gene Name	EXOSC10
UniProt Synonym Gene Names	PMSCL; PMSCL2; RRP6; PM/Scl-100
UniProt Entry Name	EXOSX_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/exosc10-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291450>