

Rabbit GPR25 Polyclonal Antibody | anti-GPR25 antibody

GPR25 Polyclonal Antibody

CATALOG #	HOST	APPLICATIONS
AAA291480	Rabbit	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry

PRODUCT SYNONYMS

GPR25; Polyclonal Antibody; GPR25 Polyclonal Antibody; Probable G-protein coupled receptor 25; anti-GPR25 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-GPR25 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Tissue/cell: human gastric carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min;Incubation: Anti-GPR25 Polyclonal Antibody, Unconjugated(bs-13526R) 1:400, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



IHC (Immunohistochemistry)

Tissue/cell: human meningioma tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min;Incubation: Anti-GPR25 Polyclonal Antibody, Unconjugated(bs-13526R) 1:500, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



IHC (Immunohistochemistry)

Tissue/cell: human glioma tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min;Incubation: Anti-GPR25 Polyclonal Antibody, Unconjugated(bs-13526R) 1:500, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

NCBI AND UNIPROT INFORMATION

NCBI GI #	38194224
NCBI GeneID	2848
NCBI Accession #	NP_005289.2

NCBI GenBank Nucleotide #	NM_005298.3
UniProt Accession #	O00155; A0AVJ5
Molecular Weight	38,779 Da
NCBI Official Full Name	probable G-protein coupled receptor 25
NCBI Official Synonym Full Names	G protein-coupled receptor 25
NCBI Official Symbol	GPR25
NCBI Protein Information	probable G-protein coupled receptor 25
UniProt Protein Name	Probable G-protein coupled receptor 25
UniProt Gene Name	GPR25
UniProt Entry Name	GPR25_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/gpr25-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-elisa/291480>