

Rabbit anti-Pig, Cow CED6 Polyclonal Antibody | anti-CED6 antibody

CED6 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291489	Rabbit	Pig, Cow	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry


PRODUCT SYNONYMS

CED6; Polyclonal Antibody; CED6 Polyclonal Antibody; PTB domain-containing engulfment adapter protein 1; Cell death protein 6 homolog; PTB domain adapter protein CED-6; Protein GULP; GULP1; GULP; anti-CED6 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Pig, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-CED6 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (mouse pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (CED6) Polyclonal Antibody, Unconjugated (bs-13626R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (mouse pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (CED6) Polyclonal Antibody, Unconjugated (bs-13626R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (CED6) Polyclonal Antibody, Unconjugated (bs-13626R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	7705318
NCBI GeneID	51454

NCBI Accession #	NP_057399.1
NCBI GenBank Nucleotide #	NM_016315.4
UniProt Accession #	Q9UBP9; Q53PC1; Q53RF3; Q9BVL3; B2RB51; B4DQ40; B8ZZ72; D3DPH1; E9PB86
Molecular Weight	19,423 Da
NCBI Official Full Name	PTB domain-containing engulfment adapter protein 1 isoform a
NCBI Official Synonym Full Names	GULP PTB domain containing engulfment adaptor 1
NCBI Official Symbol	GULP1
NCBI Official Synonym Symbols	CED6; GULP; CED-6
NCBI Protein Information	PTB domain-containing engulfment adapter protein 1
UniProt Protein Name	PTB domain-containing engulfment adapter protein 1
UniProt Gene Name	GULP1
UniProt Synonym Gene Names	CED6; GULP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ced6-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-elisa/291489>