

Rabbit TIFA Polyclonal Antibody | anti-TIFA antibody

TIFA Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291495	Rabbit	Rat, Pig, Dog, Cow	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry


PRODUCT SYNONYMS

TIFA; Polyclonal Antibody; TIFA Polyclonal Antibody; TRAF-interacting protein with FHA domain-containing protein A; Putative MAPK-activating protein PM14; Putative NF-kappa-B-activating protein 20; TRAF2-binding protein; anti-TIFA antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Pig, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-TIFA antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry)
Notes	ECO:0000303 PubMed:12566447, ECO:0000312 HGNC: HGNC:19075
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (TIFA) Polyclonal Antibody, Unconjugated (bs-13669R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



FCM/FACS (Flow Cytometry)

Blank control: Mouse spleen. Primary Antibody (green line): Rabbit Anti-TIFA antibody (bs-13669R) Dilution: 2 ug /10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG .Secondary Antibody : Goat anti-rabbit IgG-AF488 Dilution: 1 ug /test. Protocol The cells were fixed with 70% ethanol (10min at room temperature) and then were incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

NCBI AND UNIPROT INFORMATION

NCBI GI #	33563370
NCBI GeneID	92610
NCBI Accession #	NP_443096.1
NCBI GenBank Nucleotide #	NM_052864.2
UniProt Accession #	Q96CG3
Molecular Weight	21,445 Da
NCBI Official Full Name	TRAF-interacting protein with FHA domain-containing protein A

NCBI Official Synonym Full Names	TRAF-interacting protein with forkhead-associated domain
NCBI Official Symbol	TIFA
NCBI Official Synonym Symbols	T2BP; T6BP; TIFAA
NCBI Protein Information	TRAF-interacting protein with FHA domain-containing protein A; TRAF-interacting protein with a forkhead-associated domain; TRAF2 binding protein; TRAF2-binding protein; TRAF6 binding protein; putative MAPK-activating protein PM14; putative NF-kappa-B-acti
UniProt Protein Name	TRAF-interacting protein with FHA domain-containing protein A
UniProt Gene Name	TIFA
UniProt Synonym Gene Names	T2BP
UniProt Entry Name	TIFA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tifa-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-elisa/291495>