

Rabbit LXRalpha Polyclonal Antibody | anti-LXRalpha antibody

LXRalpha Polyclonal Antibody

CATALOG # AAA29155	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

LXRalpha; Polyclonal Antibody; LXRalpha Polyclonal Antibody; NR1H3; LXRA; Oxysterols receptor LXR-alpha; Liver X receptor alpha; Nuclear receptor subfamily 1 group H member 3; anti-LXRalpha antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-LXRalpha antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-LXRalpha antibody

Nuclear receptor that exhibits a ligand-dependent transcriptional activation activity (PubMed:25661920). Interaction with retinoic acid receptor (RXR) shifts RXR from its role as a silent DNA-binding partner to an active ligand-binding subunit in mediating retinoid responses through target genes defined by LXRES (By similarity). LXRES are DR4-type response elements characterized by direct repeats of two similar hexanucleotide half-sites spaced by four nucleotides (By similarity). Plays an important role in the regulation of cholesterol homeostasis, regulating cholesterol uptake through MYLIP-dependent ubiquitination of LDLR, VLDLR and LRP8 (PubMed:19481530). Interplays functionally with RORA for the regulation of genes involved in liver metabolism (By similarity).

NCBI AND UNIPROT INFORMATION

NCBI GI #	23503089
NCBI GeneID	10062
UniProt Accession #	Q13133; Q8IW13; Q96H87; A8K3J9; D3DQR1
Molecular Weight	447
NCBI Official Full Name	Oxysterols receptor LXR-alpha
NCBI Official Synonym Full Names	nuclear receptor subfamily 1, group H, member 3
NCBI Official Symbol	NR1H3
NCBI Official Synonym Symbols	LXRA; LXR-a; RLD-1
NCBI Protein Information	oxysterols receptor LXR-alpha; liver X nuclear receptor alpha variant 1
UniProt Protein Name	Oxysterols receptor LXR-alpha
UniProt Gene Name	NR1H3
UniProt Synonym Gene Names	LXRA
UniProt Entry Name	NR1H3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/lxralpha-antibody-western-blot-immunohistochemistry-elisa/29155>