

Rabbit anti-Rat, Cow DOC2B Polyclonal Antibody | anti-DOC2B antibody

DOC2B Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291558	Rabbit	Rat, Cow	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

DOC2B; Polyclonal Antibody; DOC2B Polyclonal Antibody; Double C2-like domain-containing protein beta; Doc2-beta; anti-DOC2B antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-DOC2B antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (rat kidney tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (DOC2B) Polyclonal Antibody, Unconjugated (bs-14402R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (DOC2B) Polyclonal Antibody, Unconjugated (bs-14402R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (DOC2B) Polyclonal Antibody, Unconjugated (bs-14402R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	295054132
NCBI GeneID	8447

NCBI Accession #	NP_003576.2
NCBI GenBank Nucleotide #	NM_003585.4
UniProt Accession #	Q14184
Molecular Weight	45,949 Da
NCBI Official Full Name	double C2-like domain-containing protein beta
NCBI Official Synonym Full Names	double C2-like domains, beta
NCBI Official Symbol	DOC2B
NCBI Official Synonym Symbols	DOC2BL
NCBI Protein Information	double C2-like domain-containing protein beta; doc2-beta
UniProt Protein Name	Double C2-like domain-containing protein beta
UniProt Gene Name	DOC2B
UniProt Synonym Gene Names	DOC2BL; Doc2-beta
UniProt Entry Name	DOC2B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/doc2b-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291558>