Rabbit anti-Human TNF-alpha Polyclonal Antibody | anti-TNF-alpha antibody

TNF-alpha Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA29158	**Rabbit**	**Human**	**Western Blot, Immunohistochemistry**

PRODUCT SYNONYMS

TNF-alpha; Polyclonal Antibody; TNF-alpha Polyclonal Antibody; TNF; TNFA; TNFSF2; Tumor necrosis factor; Cachectin; Tumor necrosis factor ligand superfamily member 2; TNF-a; anti-TNF-alpha antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-TNF-alpha antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)
Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistchemistry)
Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry)
Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)
Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)
Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)
Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-TNF-alpha antibody	Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. It is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia, Under certain conditions it can stimulate cell proliferation and induce cell differentiation. Impairs regulatory T-cells (Treg) function in individuals with rheumatoid arthritis via FOXP3 dephosphorylation. Upregulates the expression of protein phosphatase 1 (PP1), which dephosphorylates the key 'Ser-418' residue of FOXP3, thereby inactivating FOXP3 and rendering Treg cells functionally defective (PubMed:16829952, PubMed:23396208). Induces insulin resistance in adipocytes via inhibition of insulin-induced IRS1 tyrosine phosphorylation and insulin-induced glucose uptake. Induces GKAP42 protein degradation in adipocytes which is partially responsible for TNF-induced insulin resistance (By similarity).

NCBI AND UNIPROT INFORMATION

NCBI GI #	25952111
NCBI GeneID	7124
NCBI Accession #	NP_000585.2
NCBI GenBank Nucleotide #	NM_000594.3
UniProt Accession #	P01375; O43647; Q9P1Q2; Q9UIV3
Molecular Weight	25,644 Da
NCBI Official Full Name	tumor necrosis factor
NCBI Official Synonym Full Names	tumor necrosis factor
NCBI Official Symbol	TNF
NCBI Official Synonym Symbols	DIF; TNFA; TNFSF2; TNF-alpha

NCBI Protein Information	tumor necrosis factor; TNF-a; cachectin; APC1 protein; TNF, monocyte-derived; TNF, macrophage-derived; tumor necrosis factor-alpha; tumor necrosis factor ligand superfamily member 2
UniProt Protein Name	Tumor necrosis factor
UniProt Gene Name	**TNF**
UniProt Synonym Gene Names	TNFA; TNFSF2; TNF-a; NTF; ICD1; ICD2
UniProt Entry Name	TNFA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/polyclonal/tnf-alpha-antibody-western-blot-immunohistochemistry-elisa/29158