

Rabbit anti-Human, Rat MK Polyclonal Antibody | anti-MK antibody

MK Polyclonal Antibody

CATALOG # AAA29166	HOST Rabbit	REACTIVITY Human, Rat	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	---------------------------------	---


PRODUCT SYNONYMS

MK; Polyclonal Antibody; MK Polyclonal Antibody; MDK; MK1; NEGF2; Midkine; Amphiregulin-associated protein; ARAP; Midgestation and kidney protein; Neurite outgrowth-promoting factor 2; Neurite outgrowth-promoting protein; anti-MK antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Applicable Applications for anti-MK antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-MK antibody

Developmentally regulated, secreted growth factor homologous to pleiotrophin (PTN), which has heparin binding activity. Binds anaplastic lymphoma kinase (ALK) which induces ALK activation and subsequent phosphorylation of the insulin receptor substrate (IRS1), followed by the activation of mitogen-activated protein kinase (MAPK) and PI3-kinase, and the induction of cell proliferation. Involved in neointima formation after arterial injury, possibly by mediating leukocyte recruitment. Also involved in early fetal adrenal gland development (By similarity).

NCBI AND UNIPROT INFORMATION

NCBI GI #	396080281
NCBI GeneID	4192
NCBI Accession #	NP_001257480.1
NCBI GenBank Nucleotide #	NM_001270551.1
UniProt Accession #	P21741; Q2LEK4; Q9UCC7
Molecular Weight	15,585 Da
NCBI Official Full Name	midkine isoform a
NCBI Official Synonym Full Names	midkine (neurite growth-promoting factor 2)
NCBI Official Symbol	MDK
NCBI Official Synonym Symbols	MK; ARAP; NEGF2
NCBI Protein Information	midkine; retinoic acid inducible factor; amphiregulin-associated protein; midgestation and kidney protein; neurite outgrowth-promoting factor 2
UniProt Protein Name	Midkine
UniProt Gene Name	MDK
UniProt Synonym Gene Names	MK1; NEGF2; MK; ARAP
UniProt Entry Name	MK_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mk-antibody-western-blot-immunohistochemistry-elisa/29166>