

Rabbit anti-Rat INT11/CPSF3L Polyclonal Antibody | anti-INT11/CPSF3L antibody

INT11/CPSF3L Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291673	Rabbit	Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

INT11/CPSF3L; Polyclonal Antibody; INT11/CPSF3L Polyclonal Antibody; Integrator complex subunit 11; Int11; 3.1.27.-; Cleavage and polyadenylation-specific factor 3-like protein; CPSF3-like protein; Protein related to CPSF subunits of 68 kDa; RC-68; INTS11; CPSF3L; RC68; anti-INT11/CPSF3L antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-INT11/CPSF3L antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (INT11/CPSF3L) Polyclonal Antibody, Unconjugated (bs-16677R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37 degree C for 20 min; Incubation: Anti-INT11/CPSF3L Polyclonal Antibody, Unconjugated (bs-16677R) 1:500, overnight at 4 degree C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Mouse embryo); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (INT11/CPSF3L) Polyclonal Antibody, Unconjugated (bs-16677R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	118572557
NCBI GeneID	54973

UniProt Accession #	Q5TA45; Q5TA46; Q5TA48; Q5TA52; Q5TA53; Q5TA54; Q7L3D7; Q96HY1; A8K5S2; B3KPR3; B4DM87; G3V1S5
Molecular Weight	600
NCBI Official Full Name	Integrator complex subunit 11
NCBI Official Synonym Full Names	cleavage and polyadenylation specific factor 3-like
NCBI Official Symbol	CPSF3L
NCBI Official Synonym Symbols	RC68; INT11; RC-68; INTS11; CPSF73L
NCBI Protein Information	integrator complex subunit 11; CPSF3-like protein; related to CPSF subunits 68 kDa; protein related to CPSF subunits of 68 kDa; cleavage and polyadenylation-specific factor 3-like protein
UniProt Protein Name	Integrator complex subunit 11
UniProt Gene Name	CPSF3L
UniProt Synonym Gene Names	INTS11; RC68; Int11; CPSF3-like protein; RC-68
UniProt Entry Name	INT11_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/int11-cpsf3l-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291673>