

Rabbit L3MBTL2 Polyclonal Antibody | anti-L3MBTL2 antibody

L3MBTL2 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291685	Rabbit	Rat, Pig, Dog, Cow	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry


PRODUCT SYNONYMS

L3MBTL2; Polyclonal Antibody; L3MBTL2 Polyclonal Antibody; Lethal (3)malignant brain tumor-like protein 2; H-I (3)mbt-like protein 2; L (3)mbt-like protein 2; anti-L3MBTL2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Rat, Pig, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-L3MBTL2 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (Human stomach cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (L3MBTL2) Polyclonal Antibody, Unconjugated (bs-16880R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (rat uterus); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (L3MBTL2) Polyclonal Antibody, Unconjugated (bs-16880R) at 1:400 overnight at 4 degree C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (L3MBTL2) Polyclonal Antibody, Unconjugated (bs-16880R) at 1:400 overnight at 4 degree C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	20149698
NCBI GeneID	83746

NCBI Accession #	NP_113676.2
NCBI GenBank Nucleotide #	NM_031488.4
UniProt Accession #	Q969R5 ; Q8TEN1 ; Q96SC4 ; Q9BQI2 ; Q9UGS4
Molecular Weight	69,264 Da
NCBI Official Full Name	lethal(3)malignant brain tumor-like protein 2
NCBI Official Synonym Full Names	l(3)mbt-like 2 (Drosophila)
NCBI Official Symbol	L3MBTL2
NCBI Official Synonym Symbols	L3MBT; H-l(3)mbt-l
NCBI Protein Information	lethal(3)malignant brain tumor-like protein 2; lethal(3)malignant brain tumor-like protein 2; H-l(3)mbt-like protein 2; L(3)mbt-like protein 2
UniProt Protein Name	Lethal(3)malignant brain tumor-like protein 2
UniProt Gene Name	L3MBTL2
UniProt Synonym Gene Names	H-l(3)mbt-like protein 2
UniProt Entry Name	LMBL2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/l3mbtl2-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-elisa/291685>