

Rabbit anti-Human MDM2 Polyclonal Antibody | anti-MDM2 antibody

MDM2 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA29169	Rabbit	Human	Western Blot, Immunohistochemistry, Immunofluorescence


PRODUCT SYNONYMS

MDM2; Polyclonal Antibody; MDM2 Polyclonal Antibody; E3 ubiquitin-protein ligase Mdm2; Double minute 2 protein; Hdm2; Oncoprotein Mdm2; p53-binding protein Mdm2; anti-MDM2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-MDM2 antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.





IHC (Immunohistochemistry)

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



Application Data

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.

IHC (Immunohistochemistry-Paraffin)

Dilution: IF: 1:50-200 Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-MDM2 antibody

E3 ubiquitin-protein ligase that mediates ubiquitination of p53/TP53, leading to its degradation by the proteasome. Inhibits p53/TP53- and p73/TP73-mediated cell cycle arrest and apoptosis by binding its transcriptional activation domain. Also acts as a ubiquitin ligase E3 toward itself and ARRB1. Permits the nuclear export of p53/TP53. Promotes proteasome-dependent ubiquitin-independent degradation of retinoblastoma RB1 protein. Inhibits DAXX-mediated apoptosis by inducing its ubiquitination and degradation. Component of the TRIM28/KAP1-MDM2-p53/TP53 complex involved in stabilizing p53/TP53. Also component of the TRIM28/KAP1-ERBB4-MDM2 complex which links growth factor and DNA damage response pathways. Mediates ubiquitination and subsequent proteasome degradation of DYRK2 in nucleus. Ubiquitinates IGF1R and SNAI1 and promotes them to proteasomal degradation (PubMed:15053880, PubMed:15195100, PubMed:15632057, PubMed:16337594, PubMed:17290220, PubMed:19098711, PubMed:19219073, PubMed:19837670, PubMed:19965871, PubMed:20173098, PubMed:20385133, PubMed:20858735, PubMed:22128911). Ubiquitinates DCX, leading to DCX degradation and reduction of the dendritic spine density of olfactory bulb granule cells (By similarity). Ubiquitinates DLG4, leading to proteasomal degradation of DLG4 which is required for AMPA receptor endocytosis (By similarity).

NCBI AND UNIPROT INFORMATION

NCBI GI #	89993689
NCBI GeneID	4193
NCBI Accession #	NP_002383.2
NCBI GenBank Nucleotide #	NM_002392.5
UniProt Accession #	Q00987; Q13226; Q13297; Q13298; Q13299; Q13300; Q13301; Q53XW0; Q71TW9; Q8WYJ1; A6NL51; A8K2S6
Molecular Weight	55,233 Da
NCBI Official Full Name	E3 ubiquitin-protein ligase Mdm2 isoform a
NCBI Official Synonym Full Names	MDM2 oncogene, E3 ubiquitin protein ligase
NCBI Official Symbol	MDM2
NCBI Official Synonym Symbols	HDMX; hdm2; ACTFS
NCBI Protein Information	E3 ubiquitin-protein ligase Mdm2; oncoprotein Mdm2; Mdm2, p53 E3 ubiquitin protein ligase homolog; double minute 2, human homolog of; p53-binding protein; Mdm2, transformed 3T3 cell double minute 2, p53 binding protein
UniProt Protein Name	E3 ubiquitin-protein ligase Mdm2
UniProt Gene Name	MDM2
UniProt Synonym Gene Names	Hdm2
UniProt Entry Name	MDM2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

