

Rabbit Ataxin 3 Polyclonal Antibody | anti-ATXN3 antibody

Ataxin 3 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291720	Rabbit	Rat, Pig, Dog, Cow	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Ataxin 3; Polyclonal Antibody; Ataxin 3 Polyclonal Antibody; Ataxin-3; 3.4.19.12; Machado-Joseph disease protein 1; Spinocerebellar ataxia type 3 protein; ATXN3; ATX3; MJD; MJD1; SCA3; anti-ATXN3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Pig, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-ATXN3 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (Ataxin 3) Polyclonal Antibody, Unconjugated (bs-17208R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (human skin cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (Ataxin 3) Polyclonal Antibody, Unconjugated (bs-17208R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (mouse stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (Ataxin 3) Polyclonal Antibody, Unconjugated (bs-17208R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	290457685
NCBI GeneID	4287
UniProt Accession #	P54252; O15284; O15285; O15286; Q8N189; Q96TC3; Q96TC4; Q9H3N0; A7LFZ5; D6RDL9; E9PB63
Molecular Weight	364
NCBI Official Full Name	Ataxin-3
NCBI Official Synonym Full Names	ataxin 3
NCBI Official Symbol	ATXN3
NCBI Official Synonym Symbols	AT3; JOS; MJD; ATX3; MJD1; SCA3
NCBI Protein Information	ataxin-3; josephin; ataxin 3 variant h; ataxin 3 variant m; ataxin 3 variant ref; olivopontocerebellar ataxia 3; Machado-Joseph disease protein 1; spinocerebellar ataxia type 3 protein; Machado-Joseph disease (spinocerebellar ataxia 3, olivopontocerebella
UniProt Protein Name	Ataxin-3
UniProt Gene Name	ATXN3
UniProt Synonym Gene Names	ATX3; MJD; MJD1; SCA3
UniProt Entry Name	ATX3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/ataxin-3-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291720>