

Rabbit LULL1/TOR1AIP2 Polyclonal Antibody | anti-LULL1/TOR1AIP2 antibody

LULL1/TOR1AIP2 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291726	Rabbit	Rat, Pig, Dog, Cow	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

LULL1/TOR1AIP2; Polyclonal Antibody; LULL1/TOR1AIP2 Polyclonal Antibody; Torsin-1A-interacting protein 2; Luminal domain-like LAP1; TOR1AIP2; IFRG15; LULL1; anti-LULL1/TOR1AIP2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Pig, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-LULL1/TOR1AIP2 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (LULL1) Polyclonal Antibody, Unconjugated (bs-17326R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




WB (Western Blot)

Dilution: Sample: Lane 1: K562 (Human) Cell Lysate at 30 ug Lane 2: SH-SY5Y (Human) Cell Lysate at 30 ug Lane 3: Hela (Human) Cell Lysate at 30 ug Lane 4: HUVEC (Human) Cell Lysate at 30 ug Lane 5: U2os (Human) Cell Lysate at 30 ug Lane 6: HepG2 (Human) Cell Lysate at 30 ug Lane 7: Liver (Mouse) Lysate at 40 ug Lane 8: Liver (Rat) Lysate at 40 ug Lane 9: Kidney (Mouse) Lysate at 40 ug Lane 10: Cerebrum (Mouse) Lysate at 40 ug Lane 11: Cerebrum (Rat) Lysate at 40 ug Primary: Anti-LULL1/TOR1AIP2 (bs-17326R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 51 kD Observed band size: 51kD



WB (Western Blot)

Sample: Small intestine (Mouse) Lysate at 40 ug Primary: Anti-LULL1/TOR1AIP2(bs-17326R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 51kD Observed band size: 56kD



WB (Western Blot)

Sample: Placenta (Mouse) Lysate at 40 ug Large intestine (Mouse) Lysate at 40 ug Primary: Anti-LULL1/TOR1AIP2(bs-17326R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 51kD Observed band size: 56kD

NCBI AND UNIPROT INFORMATION

NCBI GI #	312922356
NCBI GeneID	163590

NCBI Accession #	NP_001186189.1
NCBI GenBank Nucleotide #	NM_001199260.1
UniProt Accession #	Q8NFQ8; Q9H496
Molecular Weight	51,263 Da
NCBI Official Full Name	torsin-1A-interacting protein 2 isoform b
NCBI Official Synonym Full Names	torsin A interacting protein 2
NCBI Official Symbol	TOR1AIP2
NCBI Official Synonym Symbols	NET9; LULL1; IFRG15
NCBI Protein Information	interferon alpha responsive protein; torsin-1A-interacting protein 2; luminal domain like LAP1; luminal domain-like LAP1; interferon responsive gene 15; 15 kDa interferon-responsive protein
UniProt Protein Name	Torsin-1A-interacting protein 2
UniProt Gene Name	TOR1AIP2
UniProt Synonym Gene Names	IFRG15; LULL1
UniProt Entry Name	TOIP2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/lull1-tor1aip2-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291726>