

Rabbit RPLP0 Polyclonal Antibody | anti-RPLP0 antibody

RPLP0 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291817	Rabbit	Rat, Pig, Dog, Cow	Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

RPLP0; Polyclonal Antibody; RPLP0 Polyclonal Antibody; 60S acidic ribosomal protein P0; 60S ribosomal protein L10E; Large ribosomal subunit proteinul10; anti-RPLP0 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Rat, Pig, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-RPLP0 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




WB (Western Blot)

Dilution: Sample:K562(Human) Cell Lysate at 30 ugMolt-4(Human) Cell Lysate at 30 ugU87mg(Human) Cell Lysate at 30 ugPrimary: Anti- RPLP0 (bs-1886R) at 1/1000 dilutionSecondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilutionPredicted band size: 34 kDObserved band size: 34 kD



IHC (Immunohistochemistry)

Tissue/cell: mouse embryo tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min;Incubation: Anti-RPLP0 Polyclonal Antibody, Unconjugated(bs-1886R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



IHC (Immunohistochemistry)

Tissue/cell: mouse intestine tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min;Incubation: Anti-RPLP0 Polyclonal Antibody, Unconjugated(bs-1886R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506667
NCBI GeneID	6175

NCBI Accession #	NP_000993.1
NCBI GenBank Nucleotide #	NM_001002.3
UniProt Accession #	P05388 ; Q3B7A4 ; Q9BVK4
Molecular Weight	27,436 Da
NCBI Official Full Name	60S acidic ribosomal protein P0
NCBI Official Synonym Full Names	ribosomal protein, large, P0
NCBI Official Symbol	RPLP0
NCBI Official Synonym Symbols	P0; LP0; L10E; RPP0; PRLP0
NCBI Protein Information	60S acidic ribosomal protein P0; 60S ribosomal protein L10E; acidic ribosomal phosphoprotein P0
UniProt Protein Name	60S acidic ribosomal protein P0
UniProt Gene Name	RPLP0
UniProt Entry Name	RLA0_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/rplp0-antibody-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291817>