

Rabbit anti-Rat NFE2L1 Polyclonal Antibody | anti-NFE2L1 antibody

NFE2L1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291849	Rabbit	Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry

PRODUCT SYNONYMS

NFE2L1; Polyclonal Antibody; NFE2L1 Polyclonal Antibody; Endoplasmic reticulum membrane sensor NFE2L1; Locus control region-factor 1; LCR-F1; Nuclear factor erythroid 2-related factor 1; NF-E2-related factor 1; NFE2-related factor 1; Nuclear factor; erythroid derived 2; like 1; Protein NRF1; p120 form; Transcription factor 11; TCF-11; Transcription factor NRF1; p110 form; anti-NFE2L1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-NFE2L1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (NFE2L1) Polyclonal Antibody, Unconjugated (bs-19498R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (NFE2L1) Polyclonal Antibody, Unconjugated (bs-19498R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505379
NCBI GeneID	4779
NCBI Accession #	NP_003195.1
NCBI GenBank Nucleotide #	NM_003204.2
UniProt Accession #	Q14494; Q12877; Q96FN6; D3DTU3; D3DTU5
Molecular Weight	84,704 Da
NCBI Official Full Name	nuclear factor erythroid 2-related factor 1
NCBI Official Synonym Full Names	nuclear factor, erythroid 2-like 1

NCBI Official Symbol	NFE2L1
NCBI Official Synonym Symbols	NRF1; TCF11; LCR-F1
NCBI Protein Information	nuclear factor erythroid 2-related factor 1; TCF-11; NFE2-related factor 1; NF-E2-related factor 1; transcription factor HBZ17; transcription factor LCR-F1; locus control region-factor 1; nuclear factor (erythroid-derived 2)-like 1; nuclear factor, erythr
UniProt Protein Name	Nuclear factor erythroid 2-related factor 1
UniProt Gene Name	NFE2L1
UniProt Synonym Gene Names	HBZ17; NRF1; TCF11; NF-E2-related factor 1; NFE2-related factor 1; TCF-11
UniProt Entry Name	NF2L1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/nfe2l1-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-elisa/291849>