

Rabbit SMCHD1 Polyclonal Antibody | anti-SMCHD1 antibody

SMCHD1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291873	Rabbit	Rat, Pig, Dog, Cow	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry


PRODUCT SYNONYMS

SMCHD1; Polyclonal Antibody; SMCHD1 Polyclonal Antibody; Structural maintenance of chromosomes flexible hinge domain-containing protein 1; 3.6.1.-; anti-SMCHD1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Pig, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-SMCHD1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (SMCHD1) Polyclonal Antibody, Unconjugated (bs-19929R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (SMCHD1) Polyclonal Antibody, Unconjugated (bs-19929R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (SMCHD1) Polyclonal Antibody, Unconjugated (bs-19929R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	148839305
NCBI GeneID	23347

NCBI Accession #	NP_056110.2
NCBI GenBank Nucleotide #	NM_015295.2
UniProt Accession #	A6NHR9; O75141; Q6AHX6; Q6ZTQ8; Q9H6Q2; Q9UG39
Molecular Weight	226,374 Da
NCBI Official Full Name	structural maintenance of chromosomes flexible hinge domain-containing protein 1
NCBI Official Synonym Full Names	structural maintenance of chromosomes flexible hinge domain containing 1
NCBI Official Symbol	SMCHD1
NCBI Protein Information	structural maintenance of chromosomes flexible hinge domain-containing protein 1; SMC hinge domain-containing protein 1
UniProt Protein Name	Structural maintenance of chromosomes flexible hinge domain-containing protein 1
UniProt Gene Name	SMCHD1
UniProt Synonym Gene Names	KIAA0650
UniProt Entry Name	SMHD1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/smchd1-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-elisa/291873>