

Rabbit anti-Rat, Dog Anillin Polyclonal Antibody | anti-ANLN antibody

Anillin Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291895	Rabbit	Rat, Dog	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Anillin; Polyclonal Antibody; Anillin Polyclonal Antibody; ANLN; anti-ANLN antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Dog
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-ANLN antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




WB (Western Blot)

Sample: 293T Cell (Human) Lysate at 40 ug/NIH/3T3 Cell (Mouse) Lysate at 40 ug
 Primary: Anti-Anillin (bs-20482R) at 1/300 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 124 kD
 Observed band size: 124 kD




IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (Rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (Anillin) Polyclonal Antibody, Unconjugated (bs-20482R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (Anillin) Polyclonal Antibody, Unconjugated (bs-20482R) at 1:500 overnight at 4 degree C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (Anillin) Polyclonal Antibody, Unconjugated (bs-20482R) at 1:500 overnight at 4 degree C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	546231404
NCBI GeneID	54443

NCBI Accession #	NP_001271231.1
NCBI GenBank Nucleotide #	NM_001284302.1
UniProt Accession #	Q9NQW6; Q5CZ78; Q6NSK5; Q9H8Y4; Q9NVN9; Q9NVP0
Molecular Weight	124,199 Da
NCBI Official Full Name	actin-binding protein anillin isoform 3
NCBI Official Synonym Full Names	anillin, actin binding protein
NCBI Official Symbol	ANLN
NCBI Official Synonym Symbols	scra; Scraps
NCBI Protein Information	actin-binding protein anillin
UniProt Protein Name	Actin-binding protein anillin
UniProt Gene Name	ANLN
UniProt Entry Name	ANLN_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/anillin-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291895>