

Rabbit Spindly Polyclonal Antibody | anti-SPDL1 antibody

Spindly Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291939	Rabbit	Rat, Pig, Dog, Cow	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry


PRODUCT SYNONYMS

Spindly; Polyclonal Antibody; Spindly Polyclonal Antibody; Protein Spindly; hSpindly; Arsenite-related gene 1 protein; Coiled-coil domain-containing protein 99; Rhabdomyosarcoma antigen MU-RMS-40.4A; Spindle apparatus coiled-coil domain-containing protein 1; SPDL1; anti-SPDL1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Pig, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-SPDL1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry)
Notes	ECO:0000255 HAMAP-Rule: MF_03041
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min;Incubation: Anti-Spindly Polyclonal Antibody, Unconjugated(bs-2321R) 1:500, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



IHC (Immunohistochemistry)

Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min;Incubation: Anti-Spindly Polyclonal Antibody, Unconjugated(bs-2321R) 1:500, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

NCBI AND UNIPROT INFORMATION

NCBI GI #	157388975
NCBI GeneID	54908
NCBI Accession #	NP_060255.3
NCBI GenBank Nucleotide #	NM_017785.4
UniProt Accession #	Q96EA4; Q8TEC8; Q9HD44; Q9NX97; C9JS47
Molecular Weight	70,172 Da
NCBI Official Full Name	protein Spindly

NCBI Official Synonym Full Names	spindle apparatus coiled-coil protein 1
NCBI Official Symbol	SPDL1
NCBI Official Synonym Symbols	CCDC99
NCBI Protein Information	protein Spindly; hSpindly; spindly homolog; arsenite-related gene 1 protein; coiled-coil domain containing 99; rhabdomyosarcoma antigen MU-RMS-40.4A; coiled-coil domain-containing protein 99; rrhabdomyosarcoma antigen protein MU-RMS-40.4A; spindle apparat
UniProt Protein Name	Protein Spindly
UniProt Gene Name	SPDL1
UniProt Synonym Gene Names	CCDC99; hSpindly
UniProt Entry Name	SPDLY_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/spindly-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-elisa/291939>