

Rabbit TMEFF2 Polyclonal Antibody | anti-TMEFF2 antibody

TMEFF2 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291966	Rabbit	Rat, Pig, Dog, Cow	Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

TMEFF2; Polyclonal Antibody; TMEFF2 Polyclonal Antibody; Tomoregulin-2; TR-2; Hyperplastic polyposis protein 1; Transmembrane protein with EGF-like and two follistatin-like domains; HPP1; TENB2; TPEF; anti-TMEFF2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Rat, Pig, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-TMEFF2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




WB (Western Blot)

Dilution: Sample:Cerebrum (Rat) Lysate at 40 ugPrimary: Anti-TMEFF2 (bs-3849R) at 1/1000
 dilutionSecondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilutionPredicted band size:
 37 kDObserved band size: 48 kD



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (human liver carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (TMEFF2) Polyclonal Antibody, Unconjugated (bs-3849R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (rat spinal tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (TMEFF2) Polyclonal Antibody, Unconjugated (bs-3849R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	762205501
NCBI GeneID	23671

NCBI Accession #	NP_001292074.1
NCBI GenBank Nucleotide #	NM_001305145.1
UniProt Accession #	Q9UIK5; Q2FA44; Q4ZFW4; Q53H90; Q53RE1; Q8N2R5; Q9NR15; Q9NSS5; Q9P2Y9; Q9UK65
Molecular Weight	19,472 Da
NCBI Official Full Name	tomoregulin-2 isoform 3
NCBI Official Synonym Full Names	transmembrane protein with EGF-like and two follistatin-like domains 2
NCBI Official Symbol	TMEFF2
NCBI Official Synonym Symbols	TR; HPP1; TPEF; TR-2; TENB2; CT120.2
NCBI Protein Information	tomoregulin-2; cancer/testis antigen family 120, member 2; hyperplastic polyposis protein 1; transmembrane protein TENB2
UniProt Protein Name	Tomoregulin-2
UniProt Gene Name	TMEFF2
UniProt Synonym Gene Names	HPP1; TENB2; TPEF; TR-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tmeff2-antibody-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291966>