

Rabbit TECK Polyclonal Antibody | anti-TECK antibody

TECK Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA292002	Rabbit	Rat, Pig, Dog, Cow	Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

TECK; Polyclonal Antibody; TECK Polyclonal Antibody; C-C motif chemokine 25; Chemokine TECK; Small-inducible cytokine A25; Thymus-expressed chemokine; CCL25; SCYA25; anti-TECK antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Pig, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-TECK antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IF (Immunofluorescence)

Dilution: Paraformaldehyde-fixed, paraffin embedded (rat thymus); Antigen retrieval by boiling in sodium citrate buffer (pH6) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (TECK) Polyclonal Antibody, Unconjugated (bs-5093R) at 1:400 overnight at 4 degree C, followed by a conjugated secondary (bs-0295G-Cy3) at [1:500] for 90 minutes and DAPI staining of the nuclei.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (rat thymus); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (TECK) Polyclonal Antibody, Unconjugated (bs-5093R) at 1:400 overnight at 4 degree C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



IF (Immunofluorescence)

Paraformaldehyde-fixed, paraffin embedded (rat small intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (TECK) Polyclonal Antibody, Unconjugated (bs-5093R) at 1:400 overnight at 4 degree C, followed by a conjugated secondary (bs-0295G-Cy3) at [1:500] for 90 minutes and DAPI staining of the nuclei.

NCBI AND UNIPROT INFORMATION

NCBI GI #	212286066
NCBI GeneID	6370
UniProt Accession #	O15444; Q96KJ7; A1L4J4; A6NI52; A8K9E7; B5MCA5
Molecular Weight	150
NCBI Official Full Name	C-C motif chemokine 25
NCBI Official Synonym Full Names	chemokine (C-C motif) ligand 25
NCBI Official Symbol	CCL25
NCBI Official Synonym Symbols	TECK; Ckb15; SCYA25
NCBI Protein Information	C-C motif chemokine 25; TECKvar; Ck beta-15; chemokine TECK; thymus expressed chemokine; thymus-expressed chemokine; small-inducible cytokine A25; small inducible cytokine subfamily A (Cys-Cys), member 25
UniProt Protein Name	C-C motif chemokine 25
UniProt Gene Name	CCL25
UniProt Synonym Gene Names	SCYA25; TECK
UniProt Entry Name	CCL25_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/teck-antibody-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/292002>