

Rabbit anti-Human GGT1 Polyclonal Antibody | anti-GGT1 antibody

GGT1 Polyclonal Antibody

CATALOG # AAA29201	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

GGT1; Polyclonal Antibody; GGT1 Polyclonal Antibody; GGT; Gamma-glutamyltranspeptidase 1; GGT 1; Gamma-glutamyltransferase 1; Glutathione hydrolase 1; Leukotriene-C4 hydrolase; CD224; anti-GGT1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-GGT1 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-GGT1 antibody

Cleaves the gamma-glutamyl bond of extracellular glutathione (gamma-Glu-Cys-Gly), glutathione conjugates, and other gamma-glutamyl compounds. The metabolism of glutathione releases free glutamate and the dipeptide cysteinyl-glycine, which is hydrolyzed to cysteine and glycine by dipeptidases. In the presence of high concentrations of dipeptides and some amino acids, can also catalyze a transpeptidation reaction, transferring the gamma-glutamyl moiety to an acceptor amino acid to form a new gamma-glutamyl compound. Initiates extracellular glutathione (GSH) breakdown, provides cells with a local cysteine supply and contributes to maintain intracellular GSH level. It is part of the cell antioxidant defense mechanism. Isoform 3 seems to be inactive.

NCBI AND UNIPROT INFORMATION

NCBI GI #	47678519
NCBI GeneID	2678
UniProt Accession #	P19440; Q08247; Q14404; Q8TBS1; Q9UMK1
Molecular Weight	61,410 Da
NCBI Official Full Name	GGT1
NCBI Official Synonym Full Names	gamma-glutamyltransferase 1
NCBI Official Symbol	GGT1
NCBI Official Synonym Symbols	GGT; GTG; CD224; GGT 1; D22S672; D22S732

NCBI Protein Information	gamma-glutamyltranspeptidase 1; glutamyl transpeptidase; glutathione hydrolase 1; leukotriene-C4 hydrolase; gamma-glutamyl transpeptidase
UniProt Protein Name	Gamma-glutamyltranspeptidase 1
UniProt Gene Name	GGT1
UniProt Synonym Gene Names	GGT; GGT 1
UniProt Entry Name	GGT1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ggt1-antibody-western-blot-immunohistochemistry-elisa/29201>