

Rabbit anti-Rat MAS1L Polyclonal Antibody | anti-MAS1L antibody

MAS1L Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA292018	Rabbit	Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry


PRODUCT SYNONYMS

MAS1L; Polyclonal Antibody; MAS1L Polyclonal Antibody; Mas-related G-protein coupled receptor MRG; MAS-R; MAS1-like; MRG; anti-MAS1L antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-MAS1L antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (human breast carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (MAS1L) Polyclonal Antibody, Unconjugated (bs-5926R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37 degree C for 20 min; Incubation: Anti-MAS1L Polyclonal Antibody, Unconjugated (bs-5926R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



IHC (Immunohistochemistry)
 Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min;Incubation: Anti-MAS1L Polyclonal Antibody, Unconjugated(bs-5926R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

NCBI AND UNIPROT INFORMATION

NCBI GI #	16418463
NCBI GeneID	116511
NCBI Accession #	NP_443199.1
NCBI GenBank Nucleotide #	NM_052967.1
UniProt Accession #	P35410; Q5SUN5
Molecular Weight	42,411 Da
NCBI Official Full Name	mas-related G-protein coupled receptor MRG
NCBI Official Synonym Full Names	MAS1 proto-oncogene like, G protein-coupled receptor
NCBI Official Symbol	MAS1L
NCBI Official Synonym Symbols	MRG; MAS-L; dJ994E9.2
NCBI Protein Information	mas-related G-protein coupled receptor MRG; MAS-R; MAS1-like; MAS1 oncogene-like; mas-related G protein-coupled MRG
UniProt Protein Name	Mas-related G-protein coupled receptor MRG
UniProt Gene Name	MAS1L
UniProt Synonym Gene Names	MRG; MAS-R
UniProt Entry Name	MAS1L_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/mas1l-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-elisa/292018>