

Rabbit anti-Rat, Dog Alpha 2 antiplasmin Polyclonal Antibody | anti-SERPINF2 antibody

Alpha 2 antiplasmin Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA292032	Rabbit	Rat, Dog	Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Alpha 2 antiplasmin; Polyclonal Antibody; Alpha 2 antiplasmin Polyclonal Antibody; Alpha-2-antiplasmin; Alpha-2-AP; Alpha-2-plasmin inhibitor; Alpha-2-PI; Serpin F2; SERPINF2; AAP; PLI; anti-SERPINF2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Dog
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-SERPINF2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




WB (Western Blot)

Dilution: Sample:Liver (Mouse) Lysate at 40 ugPrimary: Anti- Alpha 2 antiplasmin (bs-6188R) at 1/1000 dilutionSecondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilutionPredicted band size: 50 kDobserved band size: 52 kD



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (human liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (Alpha 2 antiplasmin) Polyclonal Antibody, Unconjugated (bs-6188R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (Alpha 2 antiplasmin) Polyclonal Antibody, Unconjugated (bs-6188R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	115583663
NCBI GeneID	5345

NCBI Accession #	NP_000925.2
NCBI GenBank Nucleotide #	NM_000934.3
UniProt Accession #	P08697; Q8N5U7; Q9UCG2; Q9UCG3; B4E1B7
Molecular Weight	54,566 Da
NCBI Official Full Name	alpha-2-antiplasmin isoform a
NCBI Official Synonym Full Names	serpin peptidase inhibitor, clade F (alpha-2 antiplasmin, pigment epithelium derived factor), member 2
NCBI Official Symbol	SERPINF2
NCBI Official Synonym Symbols	AAP; API; PLI; A2AP; ALPHA-2-PI
NCBI Protein Information	alpha-2-antiplasmin; serpin F2; alpha-2-AP; alpha-2-plasmin inhibitor; serine (or cysteine) proteinase inhibitor, clade F (alpha-2 antiplasmin, pigment epithelium derived factor), member 2
UniProt Protein Name	Alpha-2-antiplasmin
UniProt Gene Name	SERPINF2
UniProt Synonym Gene Names	AAP; PLI; Alpha-2-AP; Alpha-2-PI
UniProt Entry Name	A2AP_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/alpha-2-antiplasmin-antibody-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/292032>