

Rabbit STK38 Polyclonal Antibody | anti-STK38 antibody

STK38 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA292034	Rabbit	Rat, Pig, Dog, Cow	Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

STK38; Polyclonal Antibody; STK38 Polyclonal Antibody; Serine/threonine-protein kinase 38; 2.7.11.1; NDR1 protein kinase; Nuclear Dbf2-related kinase 1; anti-STK38 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Pig, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-STK38 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Notes	ECO:0000312 EMBL: AAH12085.1
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

Sample:Spleen (Mouse) Lysate at 40 ugLung (Mouse) Lysate at 40 ugThymus (Mouse) Lysate at 40 ugTestis (Mouse) Lysate at 40 ugA431(Human) Cell Lysate at 30 ugPrimary: Anti-STK38 (bs-6257R) at 1/1000 dilutionSecondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilutionPredicted band size: 51 kDObserved band size: 51 kD



IHC (Immunohistochemistry)

Tissue/cell: mouse embryo tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min;Incubation: Anti-STK38 Polyclonal Antibody, Unconjugated(bs-6257R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

NCBI AND UNIPROT INFORMATION

NCBI GI #	6005814
NCBI GeneID	11329
NCBI Accession #	NP_009202.1
NCBI GenBank Nucleotide #	NM_007271.2
UniProt Accession #	Q15208; Q503A1
Molecular Weight	54,190 Da
NCBI Official Full Name	serine/threonine-protein kinase 38

NCBI Official Synonym Full Names	serine/threonine kinase 38
NCBI Official Symbol	STK38
NCBI Official Synonym Symbols	NDR; NDR1
NCBI Protein Information	serine/threonine-protein kinase 38; STK38; NDR1 protein kinase; nuclear Dbf2-related 1; nuclear Dbf2-related kinase 1; Ndr Ser/Thr kinase-like protein; serine threonine protein kinase
UniProt Protein Name	Serine/threonine-protein kinase 38
UniProt Gene Name	STK38
UniProt Entry Name	STK38_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/stk38-antibody-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/292034>