

Rabbit anti-Rat MAGEB3 Polyclonal Antibody | anti-MAGEB3 antibody

MAGEB3 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA292058	Rabbit	Rat	Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MAGEB3; Polyclonal Antibody; MAGEB3 Polyclonal Antibody; Melanoma-associated antigen B3; MAGE-B3 antigen; anti-MAGEB3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-MAGEB3 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




WB (Western Blot)

Sample: Brain (Mouse) Lysate at 40 ug
Primary: Anti- MAGEB3 (bs-6823R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 39 kD
Observed band size: 39 kD




IHC (Immunohistochemistry)

Dilution: Tissue/cell: mouse testis tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37 degree C for 20 min; Incubation: Anti-MAGEB3 Polyclonal Antibody, Unconjugated (bs-6823R) 1:500, overnight at 4 degree C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



IHC (Immunohistochemistry)

Tissue/cell: Human melanoma tumor; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37 degree C for 20 min; Incubation: Anti-MAGEB3 Polyclonal Antibody, Unconjugated (bs-6823R) 1:500, overnight at 4 degree C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



IHC (Immunohistochemistry)

Tissue/cell: Rat testis tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37 degree C for 20 min; Incubation: Anti-MAGEB3 Polyclonal Antibody, Unconjugated (bs-6823R) 1:500, overnight at 4 degree C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining

NCBI AND UNIPROT INFORMATION

NCBI GI #	29171714
NCBI GeneID	4114

NCBI Accession #	NP_002356.2
NCBI GenBank Nucleotide #	NM_002365.4
UniProt Accession #	O15480; O75861; A0AVE4; B3KQ52
Molecular Weight	39,211 Da
NCBI Official Full Name	melanoma-associated antigen B3
NCBI Official Synonym Full Names	melanoma antigen family B, 3
NCBI Official Symbol	MAGEB3
NCBI Official Synonym Symbols	CT3.5
NCBI Protein Information	melanoma-associated antigen B3; MAGE-B3 antigen; cancer/testis antigen family 3, member 5
UniProt Protein Name	Melanoma-associated antigen B3
UniProt Gene Name	MAGEB3
UniProt Entry Name	MAGB3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mageb3-antibody-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/292058>