

Rabbit BMPER Polyclonal Antibody | anti-BMPER antibody

BMPER Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA292063	Rabbit	Rat, Pig, Dog, Cow	Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

BMPER; Polyclonal Antibody; BMPER Polyclonal Antibody; BMP-binding endothelial regulator protein; Bone morphogenetic protein-binding endothelial cell precursor-derived regulator; Protein crossveinless-2; hCV2; KIAA1965; anti-BMPER antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Rat, Pig, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-BMPER antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




WB (Western Blot)

Dilution: Sample: HepG2 Cell (Human) Lysate at 30 ugPrimary: Anti-BMPER (bs-6910R) at 1/300 dilutionSecondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilutionPredicted band size: 72kDObserved band size: 75kD



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (BMPER) Polyclonal Antibody, Unconjugated (bs-6910R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (BMPER) Polyclonal Antibody, Unconjugated (bs-6910R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	37693519
NCBI GeneID	168667

NCBI Accession #	NP_597725.1
NCBI GenBank Nucleotide #	NM_133468.5
UniProt Accession #	Q8N8U9; Q8TF36; A8K1P8
Molecular Weight	75,997 Da
NCBI Official Full Name	BMP-binding endothelial regulator protein
NCBI Official Synonym Full Names	BMP binding endothelial regulator
NCBI Official Symbol	BMPER
NCBI Official Synonym Symbols	CV2; CV-2; CRIM3
NCBI Protein Information	BMP-binding endothelial regulator protein
UniProt Protein Name	BMP-binding endothelial regulator protein
UniProt Gene Name	BMPER
UniProt Synonym Gene Names	KIAA1965; hCV2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/bmper-antibody-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/292063>