

Rabbit Rad23 Polyclonal Antibody | anti-Rad23 antibody

Rad23 Polyclonal Antibody

CATALOG #	HOST	APPLICATIONS
AAA292076	Rabbit	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Rad23; Polyclonal Antibody; Rad23 Polyclonal Antibody; UV excision repair protein RAD23; RAD23; anti-Rad23 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-Rad23 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



FCM/FACS (Flow Cytometry)

Blank control: Mouse spleen. Primary Antibody (green line): Rabbit Anti-Rad23 antibody (bs-7121R) Dilution: 2 ug /10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG .Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: 1 ug /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20 degree C. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



FCM/FACS (Flow Cytometry)

Blank control: Mouse spleen. Primary Antibody (green line): Rabbit Anti-Rad23 antibody (bs-7121R) Dilution: 2 ug /10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG .Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: 1 ug /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20 degree C. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

NCBI AND UNIPROT INFORMATION

NCBI GI #	398364251
NCBI GeneID	856674
NCBI Accession #	NP_010877.3
NCBI GenBank Nucleotide #	NM_001178852.3
UniProt Accession #	P32628; D3DLL2
Molecular Weight	42,367 Da
NCBI Official Full Name	Rad23p
NCBI Official Symbol	RAD23
NCBI Protein Information	Rad23p

UniProt Protein Name	UV excision repair protein RAD23
UniProt Gene Name	RAD23

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/rad23-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/292076>