

Rabbit VEGF-C Polyclonal Antibody | anti-VEGF-C antibody

VEGF-C Polyclonal Antibody

CATALOG # AAA29208	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

VEGF-C; Polyclonal Antibody; VEGF-C Polyclonal Antibody; VEGFC; Vascular endothelial growth factor C; Flt4 ligand; Flt4-L; Vascular endothelial growth factor-related protein; VRP; anti-VEGF-C antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-VEGF-C antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES


IHC (Immunohistochemistry)


Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.


IHC (Immunohistochemistry)


Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.


IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-VEGF-C antibody

Growth factor active in angiogenesis, and endothelial cell growth, stimulating their proliferation and migration and also has effects on the permeability of blood vessels. May function in angiogenesis of the venous and lymphatic vascular systems during embryogenesis, and also in the maintenance of differentiated lymphatic endothelium in adults. Binds and activates KDR/VEGFR2 and FLT4/VEGFR3 receptors.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4885653
NCBI GeneID	7424
NCBI Accession #	NP_005420.1
NCBI GenBank Nucleotide #	NM_005429.3
UniProt Accession #	P49767 ; B2R9Q8
Molecular Weight	46,883 Da
NCBI Official Full Name	vascular endothelial growth factor C preproprotein
NCBI Official Synonym Full Names	vascular endothelial growth factor C
NCBI Official Symbol	VEGFC
NCBI Official Synonym Symbols	VRP; Flt4-L
NCBI Protein Information	vascular endothelial growth factor C; VEGF-C; FLT4 ligand DHM; vascular endothelial growth factor-related protein
UniProt Protein Name	Vascular endothelial growth factor C
UniProt Gene Name	VEGFC
UniProt Synonym Gene Names	VEGF-C; Flt4-L; VRP
UniProt Entry Name	VEGFC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/vegfc-antibody-western-blot-immunohistochemistry-elisa/29208>