

Rabbit Inhibin beta-E Polyclonal Antibody | anti-INHBE antibody

Inhibin beta-E Polyclonal Antibody

CATALOG # AAA29209	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

Inhibin beta-E; Polyclonal Antibody; Inhibin beta-E Polyclonal Antibody; INHBE; Inhibin beta E chain; Activin beta-E chain; anti-INHBE antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-INHBE antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES


IHC (Immunohistochemistry)


Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.


WB (Western Blot)


Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.


IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-INHBE antibody

Inhibins and activins inhibit and activate, respectively, the secretion of follitropin by the pituitary gland. Inhibins/activins are involved in regulating a number of diverse functions such as hypothalamic and pituitary hormone secretion, gonadal hormone secretion, germ cell development and maturation, erythroid differentiation, insulin secretion, nerve cell survival, embryonic axial development or bone growth, depending on their subunit composition. Inhibins appear to oppose the functions of activins.

NCBI AND UNIPROT INFORMATION

NCBI GI #	13899338
NCBI GeneID	83729
NCBI Accession #	NP_113667.1
NCBI GenBank Nucleotide #	NM_031479.3
UniProt Accession #	P58166
Molecular Weight	38,561 Da
NCBI Official Full Name	inhibin beta E chain
NCBI Official Synonym Full Names	inhibin, beta E
NCBI Official Symbol	INHBE
NCBI Protein Information	inhibin beta E chain; activin beta E; activin beta-E chain
UniProt Protein Name	Inhibin beta E chain
UniProt Gene Name	INHBE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/inhibin-beta-e-antibody-western-blot-immunohistochemistry-elisa/29209>