

Rabbit anti-Rat CEND1 Polyclonal Antibody | anti-CEND1 antibody

CEND1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA292097	Rabbit	Rat	Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CEND1; Polyclonal Antibody; CEND1 Polyclonal Antibody; Cell cycle exit and neuronal differentiation protein 1; BM88 antigen; BM88; anti-CEND1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-CEND1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (Mouse cerebellum); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (CEND1) Polyclonal Antibody, Unconjugated (bs-7858R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (CEND1) Polyclonal Antibody, Unconjugated (bs-7858R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (CEND1) Polyclonal Antibody, Unconjugated (bs-7858R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	30795225
NCBI GeneID	51286

NCBI Accession #	NP_057648.2
NCBI GenBank Nucleotide #	NM_016564.3
UniProt Accession #	Q8N111 ; Q9NYM6
Molecular Weight	14,954 Da
NCBI Official Full Name	cell cycle exit and neuronal differentiation protein 1
NCBI Official Synonym Full Names	cell cycle exit and neuronal differentiation 1
NCBI Official Symbol	CEND1
NCBI Official Synonym Symbols	BM88
NCBI Protein Information	cell cycle exit and neuronal differentiation protein 1; BM88 antigen
UniProt Protein Name	Cell cycle exit and neuronal differentiation protein 1
UniProt Gene Name	CEND1
UniProt Synonym Gene Names	BM88
UniProt Entry Name	CEND_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cend1-antibody-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/292097>