

Rabbit anti-Human Lymphotactin Polyclonal Antibody | anti-XCL1 antibody

Lymphotactin Polyclonal Antibody

CATALOG # AAA29212	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

Lymphotactin; Polyclonal Antibody; Lymphotactin Polyclonal Antibody; XCL1; LTN; SCYC1; ATAC; C motif chemokine 1; Cytokine SCM-1; Lymphotaxin; SCM-1-alpha; Small-inducible cytokine C1; XC chemokine ligand 1; XCL2; SCYC2; Cytokine SCM-1 beta; C motif chemokine 2; XC chemokine ligand 2; anti-XCL1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-XCL1 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.




WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-XCL1 antibody

Chemotactic activity for lymphocytes but not for monocytes or neutrophils. In thymus, mediates medullary accumulation of thymic dendritic cells and contributes to regulatory T cell development, playing a role in self-tolerance establishment.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506853
NCBI GeneID	6375
NCBI Accession #	NP_002986.1
NCBI GenBank Nucleotide #	NM_002995.2
UniProt Accession #	P47992; Q52MA8
Molecular Weight	12,517 Da
NCBI Official Full Name	lymphotactin
NCBI Official Synonym Full Names	chemokine (C motif) ligand 1
NCBI Official Symbol	XCL1
NCBI Official Synonym Symbols	LTN; ATAC; LPTN; SCM1; SCM-1; SCM1A; SCYC1; SCM-1a

NCBI Protein Information	lymphotactin; SCM-1-alpha; lymphotaxin; cytokine SCM-1; c motif chemokine 1; XC chemokine ligand 1; single cysteine motif 1a; small-inducible cytokine C1; small inducible cytokine subfamily C, member 1 (lymphotactin)
UniProt Protein Name	Lymphotactin
UniProt Gene Name	XCL1
UniProt Synonym Gene Names	LTN; SCYC1
UniProt Entry Name	XCL1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/lymphotactin-antibody-western-blot-immunohistochemistry-elisa/29212>