

Rabbit anti-Rat CAF1p55/VPRBP Polyclonal Antibody | anti-CAF1p55/VPRBP antibody

CAF1p55/VPRBP Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA292192	Rabbit	Rat	Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CAF1p55/VPRBP; Polyclonal Antibody; CAF1p55/VPRBP Polyclonal Antibody; DDB1- and CUL4-associated factor 1; HIV-1 Vpr-binding protein; VprBP; Serine/threonine-protein kinase VPRBP; 2.7.11.1; Vpr-interacting protein; DCAF1; KIAA0800; RIP; VPRBP; anti-CAF1p55/VPRBP antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-CAF1p55/VPRBP antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Notes	ECO:0000312 HGNC: HGNC:30911
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



FCM/FACS (Flow Cytometry)

Blank control (Black line): Molt4 (Black). Primary Antibody (green line): Rabbit Anti-CAF1p55_VPRBP antibody (bs-9393R) Dilution: 1 ug /10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG .Secondary Antibody (white blue line): Goat anti-rabbit IgG-AF647 Dilution: 1 ug /test.Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (CAF1p55/VPRBP) Polyclonal Antibody, Unconjugated (bs-9393R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

NCBI AND UNIPROT INFORMATION

NCBI GI #	284813529
NCBI GeneID	9730
NCBI Accession #	NP_001165375.1
NCBI GenBank Nucleotide #	NM_001171904.1
UniProt Accession #	Q9Y4B6; Q2YD74; Q8TBD9; Q9HCA1; Q9UG37
Molecular Weight	118,235 Da

NCBI Official Full Name	DDB1- and CUL4-associated factor 1 isoform 2
NCBI Official Synonym Full Names	DDB1 and CUL4 associated factor 1
NCBI Official Symbol	DCAF1
NCBI Official Synonym Symbols	RIP; VPRBP
NCBI Protein Information	DDB1- and CUL4-associated factor 1; protein VPRBP
UniProt Protein Name	DDB1- and CUL4-associated factor 1
UniProt Gene Name	DCAF1
UniProt Synonym Gene Names	VprBP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/caf1p55-vprbp-antibody-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/292192>