

Rabbit anti-Human TRAIL Polyclonal Antibody | anti-TRAIL antibody

TRAIL Polyclonal Antibody

CATALOG # AAA29280	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

TRAIL; Polyclonal Antibody; TRAIL Polyclonal Antibody; TNFSF10; APO2L; Tumor necrosis factor ligand superfamily member 10; Apo-2 ligand; Apo-2L; TNF-related apoptosis-inducing ligand; Protein TRAIL; CD253; anti-TRAIL antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-TRAIL antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES





IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.


Application Data

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-TRAIL antibody

Cytokine that binds to TNFRSF10A/TRAILR1, TNFRSF10B/TRAILR2, TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4 and possibly also to TNFRSF11B/OPG (PubMed:10549288). Induces apoptosis. Its activity may be modulated by binding to the decoy receptors TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4 and TNFRSF11B/OPG that cannot induce apoptosis.

NCBI AND UNIPROT INFORMATION

NCBI GI #	300193032
NCBI GeneID	8743
NCBI Accession #	NP_001177871.1
NCBI GenBank Nucleotide #	NM_001190942.1
UniProt Accession #	P50591; A1Y9B3
Molecular Weight	32,509 Da
NCBI Official Full Name	tumor necrosis factor ligand superfamily member 10 isoform 2
NCBI Official Synonym Full Names	tumor necrosis factor (ligand) superfamily, member 10
NCBI Official Symbol	TNFSF10
NCBI Official Synonym Symbols	TL2; APO2L; CD253; TRAIL; Apo-2L

NCBI Protein Information	tumor necrosis factor ligand superfamily member 10; Apo-2 ligand; TNF-related apoptosis inducing ligand TRAIL; tumor necrosis factor (ligand) family, member 10; chemokine tumor necrosis factor ligand superfamily member 10; tumor necrosis factor apoptosis-
UniProt Protein Name	Tumor necrosis factor ligand superfamily member 10
UniProt Gene Name	TNFSF10
UniProt Synonym Gene Names	APO2L; TRAIL; Apo-2L
UniProt Entry Name	TNF10_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/trail-antibody-western-blot-immunohistochemistry-elisa/29280>