

Rabbit anti-Human CD19 Polyclonal Antibody | anti-CD19 antibody

CD19 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA29305	Rabbit	Human	Western Blot, Immunohistochemistry


PRODUCT SYNONYMS

CD19; Polyclonal Antibody; CD19 Polyclonal Antibody; B-lymphocyte antigen CD19; B-lymphocyte surface antigen B4; Differentiation antigen CD19; T-cell surface antigen Leu-12; anti-CD19 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-CD19 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.




IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-CD19 antibody

Functions as coreceptor for the B-cell antigen receptor complex (BCR) on B-lymphocytes. Decreases the threshold for activation of downstream signaling pathways and for triggering B- cell responses to antigens (PubMed:1373518, PubMed:16672701). Activates signaling pathways that lead to the activation of phosphatidylinositol 3-kinase and the mobilization of intracellular Ca (2+) stores (PubMed:9317126, PubMed:12387743, PubMed:16672701). Is not required for early steps during B cell differentiation in the blood marrow (PubMed:1373518). Required for normal levels of serum immunoglobulins, and for production of high-affinity antibodies in response to antigen challenge (PubMed:12387743, PubMed:16672701).

NCBI AND UNIPROT INFORMATION

NCBI GI #	296010921
NCBI GeneID	930
NCBI Accession #	NP_001171569.1
NCBI GenBank Nucleotide #	NM_001178098.1
UniProt Accession #	P15391; Q96S68; Q9BRD6; A0N0P9
Molecular Weight	61,128 Da
NCBI Official Full Name	B-lymphocyte antigen CD19 isoform 1
NCBI Official Synonym Full Names	CD19 molecule
NCBI Official Symbol	CD19
NCBI Official Synonym Symbols	B4; CVID3

NCBI Protein Information	B-lymphocyte antigen CD19; differentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte surface antigen B4
UniProt Protein Name	B-lymphocyte antigen CD19
UniProt Gene Name	CD19
UniProt Entry Name	CD19_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd19-antibody-western-blot-immunohistochemistry-elisa/29305>