

Mouse anti-Human CD163 Monoclonal Antibody | anti-CD163 antibody

Anti-CD163 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA293405	Mouse	Human	Immunohistochemistry


PRODUCT SYNONYMS

CD163; Monoclonal Antibody; Anti-CD163 Antibody; M130; Scavenger receptor cysteine-rich type 1 protein M130; Hemoglobin scavenger receptor; anti-CD163 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG1
Specificity	Recognizes endogenous levels of CD163 protein.
Form/Format	Mouse IgG1. Liquid in PBS containing 50% glycerol, 0.2% BSA and 0.09% (W/V) sodium azide.
Applicable Applications for anti-CD163 antibody	IHC (Immunohistochemistry)
Cellular Location	[Soluble CD163]: Secreted
Tissue Location	Expressed in monocytes and mature macrophages such as Kupffer cells in the liver, red pulp macrophages in the spleen, cortical macrophages in the thymus, resident bone marrow macrophages and meningeal macrophages of the central nervous system. Expressed also in blood. Isoform 1 is the lowest abundant in the blood. Isoform 2 is the lowest abundant in the liver and the spleen. Isoform 3 is the predominant isoform detected in the blood
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IF (Immunofluorescence)

Dilution: IF IH (1/100 - 1/300)Immunohistochemical analysis of CD163 staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



IHC (Immunohistochemistry)

Dilution: IH: (1/100 - 1/300)Immunohistochemical analysis of CD163 staining in human placenta formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

ADDITIONAL INFORMATION

Related Product Information for anti-CD163 antibody	Mouse monoclonal antibody to CD163 KLH-conjugated synthetic peptide encompassing a sequence within human CD163. The exact sequence is proprietary. Acute phase-regulated receptor involved in clearance and endocytosis of hemoglobin/haptoglobin complexes by macrophages and may thereby protect tissues from free hemoglobin-mediated oxidative damage. May play a role in the uptake and recycling of iron, via endocytosis of hemoglobin/haptoglobin and subsequent breakdown of heme. Binds hemoglobin/haptoglobin complexes in a calcium-dependent and pH- dependent manner. Exhibits a higher affinity for complexes of hemoglobin and multimeric haptoglobin of HP*1F phenotype than for complexes of hemoglobin and dimeric haptoglobin of HP*1S phenotype. Induces a cascade of intracellular signals that involves tyrosine kinase-dependent calcium mobilization, inositol triphosphate production and secretion of IL6 and CSF1. Isoform 3 exhibits the higher capacity for ligand endocytosis and the more pronounced surface expression when expressed in cells.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	344179110
NCBI GeneID	9332
NCBI Accession #	NP_004235.4
NCBI GenBank Nucleotide #	NM_004244.5
UniProt Accession #	Q86VB7; Q07898; Q07899; Q07900; Q07901; Q2VLH7; C9JIG2
Molecular Weight	125,451 Da
NCBI Official Full Name	scavenger receptor cysteine-rich type 1 protein M130 isoform a
NCBI Official Synonym Full Names	CD163 molecule
NCBI Official Symbol	CD163
NCBI Official Synonym Symbols	M130; MM130
NCBI Protein Information	scavenger receptor cysteine-rich type 1 protein M130; hemoglobin scavenger receptor; macrophage-associated antigen
UniProt Protein Name	Scavenger receptor cysteine-rich type 1 protein M130

UniProt Gene Name	CD163
UniProt Synonym Gene Names	M130; sCD163
UniProt Entry Name	C163A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd163-antibody-immunohistochemistry-elisa/293405>