

Rabbit Cleaved-PARP-1 Polyclonal Antibody | anti-PARP1 antibody

Cleaved-PARP-1 (G215) Polyclonal Antibody

CATALOG # AAA293444	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot
-------------------------------	-----------------------	--	-------------------------------------


PRODUCT SYNONYMS

Cleaved-PARP-1; Polyclonal Antibody; Cleaved-PARP-1 (G215) Polyclonal Antibody; PARP1; ADPRT; PPOL; Poly [ADP-ribose] polymerase 1; PARP-1; ADP-ribosyltransferase diphtheria toxin-like 1; ARTD1; NAD (+) ADP-ribosyltransferase 1; ADPRT 1; Poly[ADP-ribose] synthase 1; anti-PARP1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-PARP1 antibody	WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-PARP1 antibody	Poly-ADP-ribosyltransferase that mediates poly-ADP- ribosylation of proteins and plays a key role in DNA repair (PubMed:18172500, PubMed:19344625, PubMed:19661379, PubMed:23230272, PubMed:25043379, PubMed:26344098). Mainly mediates glutamate and aspartate ADP- ribosylation of target proteins: the ADP-D- ribosyl group of NAD (+) is transferred to the acceptor carboxyl group of glutamate and aspartate residues and further ADP-ribosyl groups are transferred to the 2'-position of the terminal adenosine moiety, building up a polymer with an average chain length of 20-30 units (PubMed:9315851, PubMed:19764761, PubMed:25043379). Mediates the poly (ADP-ribosyl)ation of a number of proteins, including itself, APLF and CHFR (PubMed:19764761). Also mediates serine ADP-ribosylation of target proteins following interaction with HPF1; HPF1 conferring serine specificity (PubMed:18172500, PubMed:19344625, PubMed:19661379, PubMed:23230272). ADP-ribosylation follows DNA damage and appears as an obligatory step in a detection/signaling pathway leading to the reparation of DNA strand breaks (PubMed:18172500, PubMed:19344625, PubMed:19661379, PubMed:23230272). In addition to base excision repair (BER) pathway, also involved in double-strand breaks (DSBs) repair: together with TIMELESS, accumulates at DNA damage sites and promotes homologous recombination repair by mediating poly-ADP- ribosylation (PubMed:30356214). In addition to proteins, also able to ADP-ribosylate DNA: catalyzes ADP- ribosylation of DNA strand break termini containing terminal phosphates and a 2'-OH group in single- and double-stranded DNA, respectively (PubMed:27471034). Required for PARP9 and DTX3L recruitment to DNA damage sites (PubMed:23230272). PARP1-dependent PARP9-DTX3L-mediated ubiquitination promotes the rapid and specific recruitment of 53BP1/TP53BP1, UIMC1/RAP80, and BRCA1 to DNA damage sites (PubMed:23230272). Acts as a regulator of transcription: positively regulates the transcription of MTUS1 and negatively regulates the transcription of MTUS2/TIP150 (PubMed:19344625). With EEF1A1 and TXK, forms a complex that acts as a T-helper 1 (Th1) cell-specific transcription factor and binds the promoter of IFN-gamma to directly regulate its transcription, and is thus involved importantly in Th1 cytokine production (PubMed:17177976). Involved in the synthesis of ATP in the nucleus, together with NMNAT1, PARG and NUDT5 (PubMed:27257257). Nuclear ATP generation is required for extensive chromatin remodeling events that are energy-consuming (PubMed:27257257).
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	156523968
NCBI GeneID	142
NCBI Accession #	NP_001609.2
NCBI GenBank Nucleotide #	NM_001618.3
UniProt Accession #	P09874; Q8IUZ9; B1ANJ4
Molecular Weight	113,084 Da
NCBI Official Full Name	poly
NCBI Official Synonym Full Names	poly (ADP-ribose) polymerase 1
NCBI Official Symbol	PARP1
NCBI Official Synonym Symbols	PARP; PPOL; ADPRT; ARTD1; ADPRT1; PARP-1; ADPRT 1; pADPRT-1
NCBI Protein Information	poly [ADP-ribose] polymerase 1; poly [ADP-ribose] polymerase 1; poly(ADP-ribose) polymerase; poly(ADP-ribose) synthetase; poly[ADP-ribose] synthase 1; poly(ADP-ribosyl)transferase; ADP-ribosyltransferase NAD(+); NAD(+) ADP-ribosyltransferase 1; poly (ADP-
UniProt Protein Name	Poly [ADP-ribose] polymerase 1
UniProt Gene Name	PARP1
UniProt Synonym Gene Names	ADPRT; PPOL; PARP-1; ARTD1; ADPRT 1
UniProt Entry Name	PARP1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.