

Rabbit NFkappaB-p65 Polyclonal Antibody | anti-NFkappaB-p65 antibody

NFkappaB-p65 (Acetyl Lys122) Polyclonal Antibody

CATALOG # AAA293501	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot
-------------------------------	-----------------------	--	-------------------------------------


PRODUCT SYNONYMS

NFkappaB-p65; Polyclonal Antibody; NFkappaB-p65 (Acetyl Lys122) Polyclonal Antibody; RELA; NFKB3; Transcription factor p65; Nuclear factor NF-kappa-B p65 subunit; Nuclear factor of kappa light polypeptide gene enhancer in B-cells 3; anti-NFkappaB-p65 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-NFkappaB-p65 antibody	WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-NFkappaB-p65 antibody	NF-kappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52. The heterodimeric RELA- NFKB1 complex appears to be most abundant one. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. The NF-kappa-B heterodimeric RELA- NFKB1 and RELA-REL complexes, for instance, function as transcriptional activators. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF- kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. The inhibitory effect of I-kappa-B on NF-kappa-B through retention in the cytoplasm is exerted primarily through the interaction with RELA. RELA shows a weak DNA-binding site which could contribute directly to DNA binding in the NF-kappa-B complex. Beside its activity as a direct transcriptional activator, it is also able to modulate promoters accessibility to transcription factors and thereby indirectly regulate gene expression. Associates with chromatin at the NF-kappa-B promoter region via association with DDX1. Essential for cytokine gene expression in T-cells (PubMed:15790681). The NF-kappa-B homodimeric RELA-RELA complex appears to be involved in invasin- mediated activation of IL-8 expression.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	223468681
NCBI GeneID	5970
NCBI Accession #	NP_001138610.1
NCBI GenBank Nucleotide #	NM_001145138.1
UniProt Accession #	Q04206; Q6GTV1; Q6SLK1
Molecular Weight	60,219 Da
NCBI Official Full Name	transcription factor p65 isoform 2
NCBI Official Synonym Full Names	v-rel reticuloendotheliosis viral oncogene homolog A (avian)
NCBI Official Symbol	RELA
NCBI Official Synonym Symbols	p65; NFKB3
NCBI Protein Information	transcription factor p65; NF-kappa-B p65delta3; nuclear factor NF-kappa-B p65 subunit; nuclear factor of kappa light polypeptide gene enhancer in B-cells 3
UniProt Protein Name	Transcription factor p65
UniProt Gene Name	RELA
UniProt Synonym Gene Names	NFKB3
UniProt Entry Name	TF65_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.