

Rabbit anti-Human CD314 Polyclonal Antibody | anti-CD314 antibody

CD314 Polyclonal Antibody

CATALOG # AAA29363	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

CD314; Polyclonal Antibody; CD314 Polyclonal Antibody; NKG2-D type II integral membrane protein (Killer cell lectin-like receptor subfamily K member 1)(NK cell receptor D)(NKG2-D-activating NK receptor)(CD antigen CD314); anti-CD314 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-CD314 antibody	IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: IHC-p 1:50-200, ELISA 1:10000-20000




IHC (Immunohistochemistry)

Dilution: IHC-p 1:50-200, ELISA 1:10000-20000




IHC (Immunohistochemistry-Paraffin)

Dilution: IHC-p 1:50-200, ELISA 1:10000-20000




IHC (Immunohistochemistry-Paraffin)

Dilution: IHC-p 1:50-200, ELISA 1:10000-20000



IHC (Immunohistochemistry-Paraffin)
Dilution: IHC-p 1:50-200, ELISA 1:10000-20000



IHC (Immunohistochemistry)
Dilution: IHC-p 1:50-200, ELISA 1:10000-20000

ADDITIONAL INFORMATION

Related Product Information for anti-CD314 antibody

Function as an activating and costimulatory receptor involved in immunosurveillance upon binding to various cellular stress-inducible ligands displayed at the surface of autologous tumor cells and virus-infected cells. Provides both stimulatory and costimulatory innate immune responses on activated killer (NK) cells, leading to cytotoxic activity. Acts as a costimulatory receptor for T-cell receptor (TCR) in CD8 (+) T-cell-mediated adaptive immune responses by amplifying T-cell activation. Stimulates perforin-mediated elimination of ligand-expressing tumor cells. Signaling involves calcium influx, culminating in the expression of TNF-alpha. Participates in NK cell-mediated bone marrow graft rejection. May play a regulatory role in differentiation and survival of NK cells. Binds to ligands belonging to various subfamilies of MHC class I-related glycoproteins including MICA, MICB, RAET1E, RAET1G, RAET1L/ULBP6, uLBP1, uLBP2, uLBP3 (ULBP2>ULBP1>ULBP3) and uLBP4.

NCBI AND UNIPROT INFORMATION

NCBI GI #	315221164
NCBI GeneID	100528032
NCBI Accession #	NP_001186734.1
NCBI GenBank Nucleotide #	NM_001199805.1
UniProt Accession #	P26718; Q9NR41; A8K7K5; A8K7P4
Molecular Weight	25,274 Da
NCBI Official Full Name	NKG2-D type II integral membrane protein
NCBI Official Symbol	KLRC4-KLRK1
NCBI Protein Information	NKG2-D type II integral membrane protein; NK cell receptor D; NKG2-D-activating NK receptor; killer cell lectin-like receptor subfamily K member 1
UniProt Protein Name	NKG2-D type II integral membrane protein
UniProt Gene Name	KLRK1
UniProt Synonym Gene Names	D12S2489E; NKG2D
UniProt Entry Name	NKG2D_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd314-antibody-immunohistochemistry-elisa/29363>