

Rabbit anti-Human 6CKine Polyclonal Antibody | anti-6CKine antibody

6CKine Polyclonal Antibody

CATALOG # AAA29370	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

6CKine; Polyclonal Antibody; 6CKine Polyclonal Antibody; C-C motif chemokine 21 (6CKine)(Beta-chemokine exodus-2)(Secondary lymphoid-tissue chemokine)(SLC)(Small-inducible cytokine A21); anti-6CKine antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-6CKine antibody	IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES




IHC (Immunohistochemistry)
Dilution: IHC-p 1:50-200, ELISA 1:10000-20000




IHC (Immunohistochemistry)
Dilution: IHC-p 1:50-200, ELISA 1:10000-20000




IHC (Immunohistochemistry)
Dilution: IHC-p 1:50-200, ELISA 1:10000-20000



IHC (Immunohistochemistry-Paraffin)
Dilution: IHC-p 1:50-200, ELISA 1:10000-20000



IHC (Immunohistochemistry-Paraffin)
Dilution: IHC-p 1:50-200, ELISA 1:10000-20000



IHC (Immunohistochemistry-Paraffin)
Dilution: IHC-p 1:50-200, ELISA 1:10000-20000

ADDITIONAL INFORMATION

Related Product Information for anti-6CKine antibody

Inhibits hemopoiesis and stimulates chemotaxis. Chemotactic in vitro for thymocytes and activated T-cells, but not for B-cells, macrophages, or neutrophils. Shows preferential activity towards naive T-cells. May play a role in mediating homing of lymphocytes to secondary lymphoid organs. Binds to atypical chemokine receptor ACKR4 and mediates the recruitment of beta-arrestin (ARRB1/2) to ACKR4.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4128129
NCBI GeneID	6366
UniProt Accession #	O00585
Molecular Weight	14,646 Da
NCBI Official Full Name	secondary lymphoid tissue chemokine
NCBI Official Synonym Full Names	chemokine (C-C motif) ligand 21
NCBI Official Symbol	CCL21
NCBI Official Synonym Symbols	ECL; SLC; CKb9; TCA4; 6CKine; SCYA21
NCBI Protein Information	C-C motif chemokine 21; beta chemokine exodus-2; secondary lymphoid tissue chemokine; Efficient Chemoattractant for Lymphocytes; small inducible cytokine subfamily A (Cys-Cys), member 21
UniProt Protein Name	C-C motif chemokine 21
UniProt Gene Name	CCL21
UniProt Synonym Gene Names	SCYA21; SLC
UniProt Entry Name	CCL21_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/6ckine-antibody-immunohistochemistry-elisa/29370>