

Rabbit CXCR4 Polyclonal Antibody | anti-CXCR4 antibody

CXCR4 Polyclonal Antibody

CATALOG # AAA293731	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

CXCR4; Polyclonal Antibody; CXCR4 Polyclonal Antibody; C-X-C chemokine receptor type 4; CXCR4; FB22; Fusin; HM89; LCR1; Leukocyte-derived seven transmembrane domain receptor; LESTR; NPYRL; Stromal cell-derived factor 1 receptor; SDF-1 receptor; CD antigen CD184; anti-CXCR4 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-CXCR4 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES


Application Data

Dilution: WB 1:1000-2000, IHC 1:100-200


Application Data

Dilution: WB 1:1000-2000, IHC 1:100-200


WB (Western Blot)

Dilution: WB 1:1000-2000, IHC 1:100-200

ADDITIONAL INFORMATION

Related Product Information for anti-CXCR4 antibody	Receptor for the C-X-C chemokine CXCL12/SDF-1 that transduces a signal by increasing intracellular calcium ion levels and enhancing MAPK1/MAPK3 activation (PubMed:28978524, PubMed:18799424, PubMed:24912431). Involved in the AKT signaling cascade (PubMed:24912431). Plays a role in regulation of cell migration, e.g. during wound healing (PubMed:28978524). Acts as a receptor for extracellular ubiquitin; leading to enhanced intracellular calcium ions and reduced cellular cAMP levels (PubMed:20228059). Binds bacterial lipopolysaccharide (LPS) et mediates LPS-induced inflammatory response, including TNF secretion by monocytes (PubMed:11276205). Involved in hematopoiesis and in cardiac ventricular septum formation. Also plays an essential role in vascularization of the gastrointestinal tract, probably by regulating vascular branching and/or remodeling processes in endothelial cells. Involved in cerebellar development. In the CNS, could mediate hippocampal- neuron survival (By similarity).
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	56790927
NCBI GeneID	7852
NCBI Accession #	NP_001008540.1
NCBI GenBank Nucleotide #	NM_001008540.1
UniProt Accession #	P61073; O60835; P30991; P56438; Q53S69; Q9BXA0; Q9UKN2; B2R5N0
Molecular Weight	39,746 Da
NCBI Official Full Name	C-X-C chemokine receptor type 4 isoform a
NCBI Official Synonym Full Names	chemokine (C-X-C motif) receptor 4
NCBI Official Symbol	CXCR4
NCBI Official Synonym Symbols	FB22; HM89; LAP3; LCR1; NPYR; WHIM; CD184; LESTR; NPY3R; NPYRL; HSY3RR; NPYY3R; D2S201E
NCBI Protein Information	C-X-C chemokine receptor type 4; fusin; CXC-R4; CXCR-4; CD184 antigen; SDF-1 receptor; neuropeptide Y receptor Y3; seven transmembrane helix receptor; stromal cell-derived factor 1 receptor; lipopolysaccharide-associated protein 3; seven-transmembrane-seg
UniProt Protein Name	C-X-C chemokine receptor type 4
UniProt Gene Name	CXCR4
UniProt Synonym Gene Names	CXC-R4; CXCR-4; LESTR
UniProt Entry Name	CXCR4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cxcr4-antibody-immunohistochemistry-western-blot-elisa/293731>